Impact of Medicare's Non-Payment Program on Hospital-Acquired Conditions

by

Caroline Pinto Thirukumaran

Submitted in Partial Fulfillment of the
Requirements for the Degree
Doctor of Philosophy

Supervised by Professor Yue Li

Department of Public Health Sciences
School of Medicine and Dentistry
University of Rochester
Rochester, New York

2016

BIOGRAPHICAL SKETCH

Caroline P. Thirukumaran was born in Mumbai, India. She received her Bachelor in Medicine and Surgery (MBBS) degree from the University of Mumbai in 2001, and her Master in Hospital Administration degree from the Tata Institute of Social Sciences in 2003. Following this, she gained considerable experience in hospital planning and operations through her work in tertiary care hospitals in the Indian subcontinent, the Middle East, and Africa. She began her doctoral studies in Health Services Research and Policy at the University of Rochester in 2010. She was awarded a Dean's Fellowship for her doctoral studies during 2010-2012; after which she was supported by the Department of Orthopaedics and Rehabilitation for her work as a Graduate Research Assistant. Caroline has received several merit awards for her academic achievements during her Bachelors and Masters programs. She has presented her research at several national and regional meetings, and has received awards and recognition, including the Best Student Abstract Award from the Health Information Technology Interest Group at the AcademyHealth Annual Research Meeting of 2013. She pursued her research in hospital-acquired conditions under the supervision of Professor Yue Li.

The following publications resulted from her work during her doctoral studies:

1. Raudenbush BL, **Thirukumaran CP**, Li Y, Sanders JO, Rubery PT, Mesfin A. Impact of a Comparative Study on the Management of Scoliosis in Duchenne Muscular Dystrophy: Are Corticosteroids Decreasing the Rate of Scoliosis Surgery in the United States?. *Spine (Phila Pa 1976)*. 2016. *In press*.

2. Shields E, **Thirukumaran C**, Thorsness R, Noyes K, Voloshin I. Patient factors influencing return to work and cumulative financial claims after clavicle fractures in workers' compensation cases. *Journal of Shoulder and Elbow Surgery*. 2016;25(7):1115-1121.
3. **Thirukumaran CP**, Raudenbush B, Li Y, Molinari R, Rubery P, Mesfin A. National Trends in the Surgical Management of Adult Lumbar Isthmic Spondylolisthesis: 1998 to 2011. *Spine*. 2016;41(6):490-501.
4. Shields E, **Thirukumaran C**, Thorsness R, Noyes K, Voloshin I. An Analysis of Adult Patient Risk Factors and Complications within 30 Days after Arthroscopic Shoulder Surgery. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2015;31(5):807-815.
5. **Thirukumaran CP**, Dolan JG, Webster PR, Panzer RJ, Friedman B. The Impact of Electronic Health Record Implementation and Use on Performance of the Surgical Care Improvement Project Measures. *Health Serv Res*. 2015;50(1):273-289. doi: 10.1111/1475-6773.12191.

ACKNOWLEDGEMENTS

The past six years have been a period of tremendous professional and personal growth for me, and I would like to acknowledge those who have wholeheartedly supported me during this journey.

I owe my deepest gratitude to my dissertation committee. I begin by thanking Dr. Yue Li, who is my advisor and mentor. Not only is Yue an outstanding researcher, he is also a gifted teacher. My interactions with him have made me a better researcher, and I thank him for this. I am also grateful to Dr. Helena Temkin-Greener, Dr. Laurent G. Glance, and Dr. Meredith B. Rosenthal – each of them has been generous with their guidance, encouragement, and time. I am grateful to them for their confidence in my abilities, and for enthusiastically rooting for me through the process.

My work as a Graduate Research Assistant in the Department of Orthopaedics began in 2012. Over the past four years, I have had the opportunity to work with, and learn from, talented surgeons. I am grateful to Dr. Judith F. Baumhauer and Dr. Paul T. Rubery for giving me this rich experience. Most of all, I am grateful to them for recognizing the value of my work and for giving me a once-in-a-lifetime opportunity to begin my independent research career in Orthopaedics.

I would also like to thank the Department of Public Health Sciences (PHS) faculty, both past and present, who through their advising, and teaching, have helped nurture my passion for health services research. The PHS staff deserves a special mention for

assisting me at important stages in the process, so that I could focus my attention on my research. A substantial part of the analysis for this dissertation was carried out using the computational facilities of the Center for Integrated Research and Computing (CIRC) – for this I would thank the CIRC scientists and staff for actively supporting this work. I am also grateful to the many friends, colleagues, and acquaintances that I have gained over the years, and who have made this academic process so enjoyable.

I feel incredibly fortunate to have had the chance to pursue my dreams and aspirations. This would not have been possible without the family that I am blessed with. My husband and two precious daughters – you are the love and joy of my life; thank you for keeping me on my feet, *always*. Mum and Dad – your love and prayers keep me going; thank you for being proud of all that I do.

ABSTRACT

Hospital-Acquired Conditions (HACs), commonly known as medical errors, are the 3rd leading cause of death in the United States; and have been linked to more than \$22 billion avoidable healthcare spending. In recent years, the federal government has introduced several payment and delivery reforms to control the occurrence of HACs. One such initiative is Medicare's Hospital-Acquired Condition – Present on Admission (HAC-POA) Program of 2008 (hereafter called Program), which does not reimburse hospitals for the costs of HACs among elderly Medicare beneficiaries. This dissertation evaluates the longitudinal impact of the Program on the likelihood of HACs in New York (NY) hospitals. The study hypothesizes that the Program was associated with fewer HACs, especially among patients admitted to hospitals at greater financial risk from the Program, as quantified by the Medicare Utilization Ratio (MUR).

The study uses the 2005-2012 NY State Inpatient Databases linked to five hospital-level databases. The key analytic strategy includes hierarchical logistic regressions to estimate difference-in-difference models with hospital random effects. The outcomes are any-or-none HAC indicators and individual HAC indicators among inpatient stays with common medical conditions and surgical procedures. The key independent variable is an interaction between the Program phase and the MUR quartile.

Among the primary outcomes, the Program was associated with a significant decline in the likelihood of (i) the all-or-none HAC indicator in the medical cohort, (ii) surgical site infections among patients undergoing spine surgery, and (iii) deep vein thrombosis and

pulmonary embolism among patients undergoing hip and knee replacements. Importantly, hospitals at an intermediate risk of financial disincentives from the Program demonstrate a significant decline in the likelihood of HACs as compared to hospitals at a low risk of disincentives. Furthermore, hospital-level variability play a small yet significant role in explaining HAC variance.

In conclusion, the Program-associated decline in the risk of certain HACs underscores the role of financial disincentives in motivating hospitals to reduce HACs. Given the differential effects of the Program across hospitals at varying financial risk, successful quality improvement practices in high-performing hospitals need to be scaled and disseminated to low-performing hospitals.

CONTRIBUTORS AND FUNDING SOURCES

This work was supervised by a dissertation committee comprising of Professor Yue Li (Advisor) and Professor Helena Temkin-Greener from the Department of Public Health Sciences, University of Rochester; Professor Laurent G. Glance from the Department of Anesthesiology, University of Rochester; and Professor Meredith B. Rosenthal from the Harvard T. H. Chan School of Public Health. The student independently completed all work for the dissertation. Professors Li and Glance provided the data for this dissertation. The student was awarded a Dean's fellowship for the first two years of the doctoral program, after which she was supported by the Department of Orthopaedics and Rehabilitation for her work as a Graduate Research Assistant.

TABLE OF CONTENTS

Biographical Sketch.....	ii
Acknowledgements	iv
Abstract.....	vi
Contributors and Funding Sources	viii
List of Tables.....	xi
List of Figures.....	xiii
List of Abbreviations.....	xiv
Chapter 1 - Background and Significance	1
1.1. Introduction	1
1.2. Motivation for the study	1
1.3. The Hospital-Acquired Condition – Present on Admission Program	4
1.4. Specific Aims and Hypotheses	9
1.5. Theoretical and Conceptual Framework.....	11
1.6. Literature Review	19
Chapter 2 - Methodology	46
2.1. Data sources.....	46
2.2. Cohort specification.....	49
2.3. Datasets and variables.....	52
2.4. Statistical analysis.....	63
2.5. Summary of key methods terms.....	73
Chapter 3 - Results	74
3.1. Descriptive statistics.....	74

3.2. Specific Aim 1	88
3.3. Specific Aim 2.1	96
3.4. Specific Aim 2.2	103
3.5. Specific Aim 2.3	113
3.6. Stratified Analysis	119
3.7. Sensitivity Analysis.....	122
Chapter 4 - Discussion	134
4.1. Findings	136
4.2. Discussion and Policy Implications.....	142
4.3. Study limitations.....	151
4.4. Conclusion	156
List of References	157
Appendix	175

LIST OF TABLES

Table 1: Hospital-Acquired Conditions evaluated in this dissertation	7
Table 2: Studies evaluating the impact of the HAC Program on HAC rates	29
Table 3: ICD-9-CM Diagnosis and Procedure codes used for cohort identification	51
Table 4: Dataset and variable linkages.....	52
Table 5: Descriptive statistics for hospitals in the medical and surgical cohorts.....	76
Table 6: Descriptive statistics for inpatient stays in the medical and surgical cohorts	78
Table 7: Unadjusted hospital-level Hospital-Acquired Condition rates in the cohort ¹	85
Table 8: Odds ratios of risk factors for the any-or-none indicator in the pre-Program phase ¹	91
Table 9: Multivariate odds ratios from hierarchical logistic regression models for the Program effects on the primary outcomes ¹	99
Table 10: Multivariate odds ratios for the primary outcomes ¹ among hospitals with differential operating profit margins	105
Table 11: Multivariate odds ratios for the primary outcomes ¹ among hospitals with differential disproportionate patient percent.....	110
Table 12: Multivariate odds ratios for the primary outcomes ¹ among hospitals with differential Electronic Health Record (EHR) capabilities	116
Table 13: Multivariate odds ratios for Hospital-Acquired Conditions in diagnosis-based strata of the medical cohort	120
Table 14: Multivariate odds ratios for primary outcomes ¹ in the short- and long-term post-Program phase.....	123

Table 15: Multivariate odds ratios for primary outcomes ¹ with redefined post-Program phases	126
Table 16: Multivariate odds ratios for primary outcomes ¹ with redefined Medicare Utilization Ratio variable	129
Table 17: Multivariate odds ratios from analysis for primary outcomes ¹ excluding hospitals with less than 20 observations a year	132
Table 18: Summary of findings for primary outcomes	139

LIST OF FIGURES

Figure 1: Change in marginal cost of quality with HAC Program	13
Figure 2: Conceptual framework for hospital response to HAC Program disincentives..	14
Figure 3: Scope of the Dissertation	18
Figure 4: Cohort identification algorithm	50
Figure 5: Lowess graphs for unadjusted mean rates for primary Hospital-Acquired Condition outcomes.....	82
Figure 6: Multivariate odds ratios from models evaluating the impact of the Hospital-Acquired Condition Program	101

LIST OF ABBREVIATIONS

AHRQ	Agency for Healthcare Research and Quality
AIR	Air Embolism
AMI	Acute Myocardial Infarction
BLD	Blood Incompatibility
BSI	Bloodstream Infection
CABG	Coronary Artery Bypass Graft
CabgSSI	Coronary Artery Bypass Graft – Surgical Site Infection
CAUTI	Catheter-Associated Urinary Tract Infection
CC	Complication / Comorbidity
CDC	Centers for Disease Control and Prevention
CDR	Clinical Data Repository
CDSS	Clinical Decision Support System
CHF	Congestive Heart Failure
CLABSI	Central Line-Associated Bloodstream Infection
COR	Conditional Odds Ratio
CMS	Centers for Medicare and Medicaid Services
DHHS	Department of Health and Human Services
DID	Difference-in-Difference
DPP	Disproportionate Patient Percent
DRA	Deficit Reduction Act
DSH	Disproportionate Share Hospital
DVT	Deep Vein Thrombosis

EBP	Evidence-based Practices
EHR	Electronic Health Record
EMAR	Electronic Medication Administration Record
EMRAM	Electronic Medical Record Adoption Model
FY	Fiscal Year
FAL	Falls and Fractures
FP	For-profit
GDP	Gross Domestic Product
GLU	Glycemic control
HAC	Hospital-Acquired Condition
HAI	Hospital-Acquired Infection
HCUP	Healthcare Cost and Utilization Project
HIMSS	Healthcare Information and Management Systems Society
HHI	Herfindahl-Hirschman index
HipDVT	Hip Replacement – Deep Vein Thrombosis
HMO	Health Maintenance Organization
HQID	Hospital Quality Incentive Demonstration
HRR	Hospital Referral Region
HSA	Hospital Service Area
ICC	Intraclass Correlation Coefficient
ICD	International Classification of Diseases
IPPS	Inpatient Prospective Payment System
KneeDVT	Knee Replacement – Deep Vein Thrombosis
LIS	Laboratory Information System

MCC	Major Complication / Comorbidity
MS-DRG	Medical Severity - Diagnosis Related Group
MUR	Medicare Utilization Ratio
NFP	Not-for-Profit
NHE	National Health Expenditure
NHSN	National Healthcare Safety Network
NSQIP	National Surgical Quality Improvement Program
NY	New York
OECD	Organisation for Economic Co-operation and Development
P4P	Pay-for-Performance
PE	Pulmonary Embolism
PMS	Pharmacy Management System
POA	Present on Admission
PPACA	Patient Protection and Affordable Care Act
QI	Quality Improvement
RIS	Radiology Information System
SpineSSI	Spine Surgery – Surgical Site Infection
US	United States
UTI	Urinary Tract Infection
VBP	Value-Based Purchasing

Chapter 1 - BACKGROUND AND SIGNIFICANCE

1.1. Introduction

There is an incontestable need to improve the quality and control the costs of health care delivered in the United States (US). Several landmark reports and peer-reviewed papers have underscored the urgency to remedy this quality-cost conundrum, and the consequences of maintaining a status quo ¹⁻⁵. Pay-for-performance programs which align financial incentives to quality metrics (the 'carrot' approach), and their mirror image of not paying for poor quality care (the 'stick' approach), are two payment strategies that have gained immense popularity in the past decade ^{6, 7}. However, rigorous empirical evaluations of these efforts are few ⁸⁻¹⁰. One such initiative that merits evaluation is the HAC-POA Program that was mandated by the Deficit Reduction Act of 2005, through which the Centers for Medicare and Medicaid Services (CMS) does not reimburse hospitals for the costs incurred in the care of HACs.

1.2. Motivation for the study

The HAC Program is one of several federal initiatives that aim to enhance the efficiency of care by targeting 'waste' that exists within the health care system. In 2011, 34% of the national health expenditure i.e. approximately \$910 billion (estimates range from \$558 billion to \$1,263 billion), were lost in activities that did not generate value ². Of the wasted resources, approximately 18% (\$163 billion) were lost due to failures of care delivery and care coordination ^{2, 11-14}. Examples of these include failure to adequately mobilize a bed-ridden stroke patient resulting in the development of pressure ulcers or failure to comply with infection control practices while inserting urinary catheters thereby

resulting in catheter-associated urinary tract infections. Importantly, while literature suggests that a large portion of these failures can be prevented with evidence-based guidelines¹⁵⁻¹⁷, an important concern is that complying with evidence-based guidelines is challenging¹⁸⁻²¹

Medicare is the single largest purchaser of health care in the US. In 2014, Medicare alone purchased 20% (\$619 billion) of the \$3 trillion spent on health care²². Over the next decade, as more Americans become eligible for Medicare, the increase in Medicare expenditure is expected to escalate by 82%, to \$1 trillion in 2021. Furthermore, inpatient hospital care consumed the largest share (42% - \$231 billion) of Medicare's resources. Thus, identifying and addressing inefficiencies within the largest sector of the largest payer is certain to contribute in enhancing the efficiency of care delivered.

According to researchers at the Dartmouth Institute for Health Policy and Clinical Practice, 30% of all Medicare spending on clinical care could be reduced without adversely influencing health outcomes²³ and a large part of this wasted spending is attributed to preventable complications that occur in hospitalized Medicare beneficiaries^{24, 25}. Several factors place elderly patients at an increased risk for these errors. The complexity of care, prolonged duration of hospitalization, multiple comorbidities, large number of medications, and impaired cognitive responses²⁶ are some of the reasons why almost one-third of Medicare beneficiaries experience adverse events related to care during their hospital stay. Importantly, 44% of these events may have been prevented by the use of evidence-based guidelines^{27, 28}. Consequently, nearly one-fifth of Medicare beneficiaries are readmitted within 30 days of their discharge from the

hospital ²⁹, and 90% of these readmissions could have been potentially prevented thereby saving \$17 billion annually ³⁰. Thus, there is a compelling case to reduce the occurrence of HACs among Medicare beneficiaries by employing innovative delivery and payment reforms.

The central aim of this dissertation is to evaluate the HAC Program's role in preventing HACs from occurring. The study uses profit-maximization theory, the Agency for Healthcare Research and Quality's (AHRQ) conceptual framework of hospital response to financial incentives, and hierarchical statistical methods to explain and test hospital response to the HAC Program.

1.3. The Hospital-Acquired Condition – Present on Admission Program

The HAC-POA Program of 2008 was mandated by the Deficit Reduction Act of 2005. According to this Act, the Secretary of the Department of Health and Human Services was required to identify HACs that met the following criteria. First, the conditions were not present in Medicare beneficiaries at the time of admission and were sustained during hospitalization because of poor quality care. Second, the HACs were high cost or high volume or both. Third, the presence of HACs would lead to the reassignment of the inpatient stay to a higher paying MS-DRG. Finally, the HACs could have been reasonably averted through the adoption of evidence-based guidelines.

The motivation for this Program can be traced back to the Institute of Medicine's landmark report "To Err is Human", which placed an unprecedented focus on the state of patient safety in American hospitals ³¹. In 2001, the National Quality Forum's list of 'never events' further underscored the need for expediting prevention of medical errors that were easily identifiable, potentially preventable, and should not occur as a consequence of care processes ³². In 2005, HealthPartners, which is a not-for-profit health maintenance organization, was the first US purchaser that did not pay providers for conditions that patients should not have acquired during the course of treatment ³³. Since then, several payment and delivery reforms have been instituted at the federal and local levels to motivate hospitals and physicians to optimize the quality of care they render. One category of payment reforms is the pay-for-performance ('carrot') approach and its mirror image of nonpayment-for-nonperformance ('stick' approach). The HAC-POA Program is an example of the latter approach whereby hospitals are not reimbursed for the cost of HAC care if Medicare beneficiaries develop HACs during their

stay. Importantly, this Program was intended to address the 'design flaw' that existed within the Inpatient Prospective Payment System (IPPS) through which, prior to the HAC Program implementation, hospitals were reimbursed an added amount for the cost of HACs through the outlier payment mechanism.

Under the IPPS, CMS reimburses hospitals a fixed predetermined amount for discharges under a particular Medical Severity – Diagnosis Related Group (MS-DRG) regardless of the length of stay or resources used. Thus, to enhance their profitability, hospitals are incentivized to minimize costs incurred during the delivery of care. However, providing high quality care for extremely sick cases can result in expenses greater than those provisioned by the MS-DRG. To enable hospitals to provide such care to these patients, Medicare has a provision termed 'outlier payments' through which hospitals are reimbursed higher amounts for extraordinarily high expenditure³⁴⁻³⁶. Prior to the HAC Program, classifying a discharge to a MS-DRG with complication/comorbidity (CC) or with major complication/comorbidity (MCC) provided the hospitals with the means to code for HACs, and resulted in higher reimbursements as compared to a MS-DRG without CC/MCC³⁷. This ability to code for complications as secondary diagnoses and to continue receiving incremental reimbursements provided hospitals with perverse incentives within the scope of the IPPS. Hospitals were able to obtain higher payments for conditions acquired because of poor quality care, and hospitals that took measures to control the occurrence of HACs encountered a decline in revenue and profits^{7, 36, 38, 39}. To address this flaw, the CMS introduced the HAC-POA Program for discharges occurring after October 1, 2008 and this is a federal program that applies to all hospitals that were reimbursed by the IPPS.

Through the HAC-POA Program, when a HAC occurs, CMS reimburses the hospital as though the particular condition or code was not present at admission. Thus, the inpatient stay is not assigned to a higher paying MS-DRG i.e. to the MS-DRG to which the case may have been assigned if the particular secondary diagnosis was present at admission. The resulting loss in revenue is intended to serve as a penalty for the hospital's failure in preventing HACs from occurring ⁴⁰. Furthermore, the hospital is not permitted to bill the patient for the difference in reimbursements ⁴¹. In contrast, a case with an HAC as a secondary diagnosis may be assigned to a higher paying MS-DRG if certain complications or comorbidities are present at admission ⁴². For example, a discharge with a primary diagnosis of intracranial hemorrhage or cerebral infarction (stroke) with MCC (MS-DRG 064), and a secondary diagnosis of Stage III pressure ulcer (ICD-9 code 707.23 coded as a MCC). If the secondary diagnosis is present at admission, hospitals are reimbursed a median amount of \$8,030.28 whereas if the secondary diagnosis is not present at admission but develops during hospital stay, the median reimbursement amount is \$5,347.98 ⁴³. The hospital thus needs to undertake a loss of \$2,682.30 for costs associated with the treatment of the pressure ulcer.

Furthermore, as a part of the Hospital Inpatient Quality Reporting Program, CMS has publicly reported hospital-level rates of 8 HACs on the Hospital Compare website since FY 2012 ⁴⁴. For public disclosure, HAC rates are calculated (as HACs per 1,000 fee-for-service discharges) using all occurrences of qualifying secondary diagnosis and POA codes, and do not exclude discharges with reassignment of MS-DRG solely due to HACs ^{40, 45}.

In this dissertation, I have examined the impact of the Program on HACs (Table 1) that were within the scope of the Program until 2012. While these HACs apply to Medicare beneficiaries admitted to hospitals across the United States that are reimbursed by IPPS, this dissertation examines the impact of the Program only among hospitals in New York State.

Table 1: Hospital-Acquired Conditions evaluated in this dissertation

Sr. No.	HAC	ICD-9-CM Codes
1	Foreign object retained after surgery	998.4, 998.7
2	Air embolism	999.1
3	Blood Incompatibility	999.60, 999.61, 999.62, 999.63, 999.69
4	Pressure ulcer (Stages III & IV)	707.23, 707.24
5	Falls and trauma	Fracture : 800-829, Dislocation: 830-839, Intracranial injury: 850-854, Crushing injury: 925-929, Burn: 940-949, Other injuries: 991-994
6	Catheter-associated urinary tract infection	996.64
7	Vascular catheter-associated infection	999.31, 999.32, 999.33
8	Manifestations of poor glycemic control	Diabetic ketoacidosis: 250.10-250.13 Nonketotic hyperosmolar coma: 250.20-250.23 Hypoglycemic coma: 251.0 Secondary diabetes with ketoacidosis: 249.10-249.11 Secondary diabetes with Hyperosmolarity: 249.20-249.21
9.1	Surgical site infection, mediastinitis, following Coronary Artery Bypass Graft (CABG)	519.2, and One of the following procedure codes: 36.10-36.19
9.2	Surgical site infection following certain orthopedic procedures (Spine, neck, shoulder, elbow)	996.67, 998.59 and One of the following procedure codes: 81.01-81.08, 81.23-81.24, 81.31-81.38, 81.83 or 81.85

Sr. No.	HAC	ICD-9-CM Codes
9.3	Surgical site infection following bariatric surgery for obesity (Laparoscopic gastric bypass, gastroenterostomy, laparoscopic gastric restrictive surgery)	Principal Diagnosis: 278.01 539.01, 539.81, 998.59, and One of the following procedure codes: 44.38, 44.39, 44.95
10	Deep vein thrombosis and pulmonary embolism following certain orthopedic procedures (Total knee replacement, hip replacement)	415.11, 415.13, 415.19, 453.40-453.42, and One of the following procedure codes: 00.85-00.87, 81.51-81.52, or 81.54

Abbreviations: ICD-9-CM – International Classification of Diseases, Ninth Revision, Clinical Modification

In fiscal year (FY) 2011, falls and trauma (16.1 per 1,000 eligible discharges), stage III and IV pressure ulcers (10.8 per 1,000 eligible discharges), and deep vein thrombosis and pulmonary embolism following certain orthopedic procedures (7.59 per 1,000 eligible discharges) were the three most common HACs among Medicare beneficiaries⁴². Incremental costs of treating HACs can range from \$1,600 for each bariatric surgery SSI to \$ 12,000 for each CABG SSIs⁴⁶, and it is estimated that CMS would save at least \$145 million annually by withholding reimbursement for incremental HAC costs⁴⁷.

1.4. Specific Aims and Hypotheses

The specific aims and hypotheses for the dissertation are presented here to give the reader a snapshot of the study objectives. The theoretical and conceptual frameworks that motivate these aims are discussed in the subsequent sections of this Chapter. The aims and hypotheses are recapitulated at relevant locations in subsequent Chapters for convenience of the reader.

This dissertation is organized into two main specific aims:

Specific Aim 1

To explain the variation in hospital-level HAC likelihood prior to the implementation of the HAC Program.

Specific Aim 2

To evaluate the impact of the HAC Program on likelihood of HACs among hospitalized Medicare beneficiaries, focusing specifically on key hospital characteristics that are likely to influence the hospital's response to the Program.

- Hypothesis 2.1: Controlling for relevant variables, the HAC Program is associated with a decline in the HAC likelihood among Medical beneficiaries admitted to NY hospitals, and that this *decline is greater* among hospitals at a risk of *higher financial disincentives from the Program*.
- Hypothesis 2.2: Controlling for relevant variables, the HAC Program is associated with a decline in the HAC likelihood among Medical beneficiaries admitted to NY hospitals, and that this *decline is greater* among hospitals at a risk of *higher financial disincentives from the Program and in better financial health*.

- Sub-hypothesis 2.2.1: Controlling for relevant variables, the HAC Program is associated with a decline in the HAC likelihood among Medical beneficiaries admitted to NY hospitals, and that this *decline is greater* among hospitals at a risk of *higher financial disincentives from the Program* and with *higher profit margins*.
- Sub-hypothesis 2.2.2: Controlling for relevant variables, the HAC Program is associated with a decline in the HAC likelihood among Medical beneficiaries admitted to NY hospitals, and that this *decline is greater* among hospitals at a risk of *higher financial disincentives from the Program* and with *lower Disproportionate Patient Percent (DPP)*.
- Hypothesis 2.3: Controlling for relevant variables, the HAC Program is associated with a decline in the HAC likelihood among Medical beneficiaries admitted to NY hospitals, and that this *decline is greater* among hospitals at a risk of *higher financial disincentives from the Program* and with *robust Electronic Health Record (EHR) capabilities*.

Each of these hypotheses is tested for relevant HACs in cohorts of medical and surgical inpatient stays.

1.5. Theoretical and Conceptual Framework

1.5.1. Theoretical Framework

I have used the economic profit maximization framework to predict the change in quality in response to disincentives of the HAC Program. The profit-maximizing objective function represents the behavior of for-profit hospitals that deliver a mix of services that maximize financial gains ⁴⁸. In contrast, the behavior of not-for-profit (NFP) hospitals is less clearly understood. Several theories, such as NFP as “for-profits in disguise”, prestige maximizers, social-welfare maximizers, and perquisite maximizers have been studied ⁴⁹. I use the NFP as “for-profits in disguise” approach to support a profit-maximizing objective function for NFP hospitals ⁵⁰. An emerging body of literature has demonstrated profitability as a part of the objective function of NFP hospitals. Arguments such as NFP hospitals maximize the profits / surplus they earn from patients reimbursed by certain payers (such as private insurance) in order to provide care to other groups of patients (such as those reimbursed by public insurance) ⁵¹ and NFP hospitals tend to balance their social commitments with financial viability have been used to explain this behavior ^{49, 52}.

Before the HAC Program:

Using the aforementioned discussion, the objective function of both for-profit and not-for-profit hospitals can be represented by a profit-maximizing behavior as follows:

$$\begin{aligned}\pi &= R - C \\ &= (P \times Q_0) - C(Q_0, q_0)\end{aligned}$$

Where,

π is the profit, which the hospital maximizes

R is the revenue of the hospital. This is composed of price P and quantity Q_0 of services provided

C is the cost of providing services by the hospital. This is a function of quantity Q_0 and quality q_0 of the services provided

The first order conditions for this equation are:

$$\frac{\partial \pi}{\partial Q_0} = P - \frac{\partial C}{\partial Q_0} = 0$$

$$\frac{\partial \pi}{\partial q_0} = - \frac{\partial C}{\partial q_0} = 0$$

Thus, from the above equation, the marginal cost of quality is equal to zero.

After the implementation of the HAC Program:

The profit maximization function of a hospital may be represented by:

$$\begin{aligned} \pi &= R - C - D \\ &= (P \times Q_1) - C(Q_1, q_1) - D(q_1) \end{aligned}$$

Where,

D are the disincentives of the HAC Program, which are a function of quality q_1 , which in turn represents the magnitude of HACs in a hospital.

The first order conditions for this equation are:

$$\frac{\partial \pi}{\partial Q_1} = P - \frac{\partial C}{\partial Q_1} = 0$$

$$\frac{\partial \pi}{\partial q_1} = - \frac{\partial C}{\partial q_1} - \frac{\partial D}{\partial q_1} = 0$$

Thus, from the above equation, the marginal cost of quality with the implementation of the HAC Program is $-\frac{\partial D}{\partial q_1}$.

Figure 1 represents the change in the marginal cost of quality with the HAC Program. Prior to the Program, the marginal cost is equal to zero. Implementation of HAC Program disincentives result in a marginal cost of $-\frac{\partial D}{\partial q_1}$. As quality q_1 increases, disincentives D will decline, assigning a negative sign to $\frac{\partial D}{\partial q_1}$, and resulting in positive marginal cost of quality with the HAC Program. This marginal cost plotted on the marginal cost curve demonstrates higher quality q_1 with the HAC Program.

Figure 1: Change in marginal cost of quality with HAC Program

1.5.2. Conceptual Model for Hospital Response to the Program

I adapt the framework (Figure 2) proposed by Dudley et al ⁵³ and Stone et al ⁴¹ to conceptually organize and explain the response of hospitals to the disincentives of the HAC Program.

Figure 2: Conceptual framework for hospital response to HAC Program disincentives

The disincentives of the HAC Program have financial and non-financial implications for hospitals, which are intended to motivate hospitals (the leadership and staff) to adopt quality-oriented behavior and evidence-based practices, thereby reducing the likelihood of HACs and consequently minimizing the risk of financial losses. The HAC Program directly influences the finances of the hospital by not reimbursing hospitals for the costs of HACs. These costs are borne by the hospital. Thus, it is in the interest of the hospitals to prevent HACs from occurring ⁵⁴. The Program may indirectly influence the financial resources of the hospital by motivating investments in quality improvement (QI) efforts for averting HACs. While considering these investments in quality, the hospital management is certain to weigh the costs and benefits resulting from the QI initiatives. The non-financial / reputational impact of the Program on hospital behavior may mainly result from the reporting of HAC rates on the Hospital Compare website. Several stakeholders, including patients and providers, may use publicly reported information while choosing between hospitals or ascertaining performance of competing hospitals. The choices made by patients have implications for the market share of the hospital, thereby incentivizing hospitals to control HACs ⁵⁵⁻⁵⁹.

Since the disincentives of the HAC Program are directed towards the hospital organization, the traits of the organization leadership and the decisions they make ^{60, 61} significantly influence the response of the hospital to the HAC Program. This emphasis on quality created by a committed leadership, a favorable work culture and a high-performing workforce are expected to facilitate structural changes such as adoption of EHRs and process changes such as integration of evidence-based practices with the work environment, which are likely to result in reduction of HACs ⁶².

Translation of evidence-based medicine into evidence-based practices is an involved process. This is moderated by several patient-level factors such as age, sex, and race; and hospital-level factors such as ownership, geographic location, and staffing ratios ⁵³,

⁶³.

1.5.3. Modeling hospital risk

Based on the theoretical and conceptual framework, it is expected that hospitals will adopt quality improvement measures that will reduce the likelihood of HACs in response to the Program. Since the Program targets the Medicare reimbursements that a hospital receives, it is likely that the sensitivity of hospitals to the Program may vary across hospitals based on the Medicare reimbursements that are at stake ⁶⁴. I have used the Medicare Utilization Ratio (MUR) to quantify this exposure of a hospital to the Program ⁶⁵. MUR is the proportion of total inpatient days for a hospital that are attributed to Medicare. Elderly Medicare patients are at a greater risk of medical errors because of larger number of comorbid conditions, frailty, and interactions of multiple caregiving teams ⁶⁶. Thus hospitals with longer staying elderly patients, and consequently with higher MUR, are at a greater risk of financial losses from the Program. Furthermore, the MUR is utilized by the CMS for calculating the annual updates to the reimbursement that hospitals receive ⁶⁵. Thus, hospitals with higher MUR can be expected to have greater sensitivity to the Program. Other measures, such as the volume of Medicare beneficiaries, and the magnitude of Medicare dollars that a hospital receives may be used to quantify the exposure or risk of a hospital to the Program. However, using a measure such as the volume of Medicare beneficiaries has certain limitations since it

does not effectively quantify the risk of HACs among the Medicare population in a hospital. As an example, a hospital with high volume of short-staying patients may have lower likelihood of HACs and hence may not need intensive quality improvement programs as compared to a hospital with lower volume but longer staying Medicare beneficiaries who are likely to be at a greater risk of HACs.

1.5.4. Integrating Economic Theory and Conceptual Model

Figure 3 integrates the conceptual model and economic theory to outline the dissertation scope. A detailed literature review that supports each of the Specific Aims is documented in Section 1.6. Specific Aim 2 will examine the overall impact of the HAC Program on HAC rates in hospitals (Hypothesis 2.1). From economic theory, I expect the Program to be associated with a decline in the likelihood of HACs. I will control for several patient- and hospital-level contextual factors that are likely to moderate the intended Program effect. The dissertation will also examine the differential effect of the Program in hospitals with varying financial health (Hypothesis 2.2). Financial health is a contextual factor that influences the association between HAC Program disincentives and the likelihood of HACs. Financial health will be represented by the profit margin of the hospital and the Disproportionate Patient Percent. Hypothesis 2.3 will examine the differential impact of the Program in hospitals with varying EHR capabilities. EHRs are structural factors that are important for facilitating quality-oriented processes and consequently resulting in decline in the likelihood of HACs.

Figure 3: Scope of the Dissertation

1.6. Literature Review

In this section, I discuss the literature that supports the formulation of the specific aims and related hypotheses.

1.6.1. Specific Aim 1

Patient and hospital characteristics are important predictors of the outcomes of care in hospitals ⁶⁷⁻⁶⁹. Identifying significant patient- and hospital-level predictors of HACs will provide important insights into the variation of HACs.

Patient-level factors

Age:

The physical and mental health status of elderly patients places them at an increased risk of developing HACs ⁷⁰⁻⁷⁵. A larger number of comorbid conditions, longer hospital stay, frailty, need for larger number of medications are some of the factors that increase the risk for elderly patients. Importantly, hospitalized patients 65 or older are at double the risk of adverse events as compared to those between 16 to 44 years of age ^{24, 66}. Falls, pressure ulcers are examples of HACs that are more common among hospitalized elderly patients ^{76, 77}.

Sex:

Male patients have been found to be at a higher risk of developing healthcare-associated CLABSIs and SSIs on account of greater bacterial colonization of the skin and other anatomical factors ⁷⁸, whereas female patients are at an increased risk of CAUTIs on account of the anatomy of the female genitourinary tract ^{79, 80}.

Race and Ethnicity:

Blacks have been found to be at a greater risk of certain nosocomial infections, and adverse drug events as compared to whites⁸¹. Potential explanations for these findings include higher number of comorbidities and longer length of hospital stay for Black patients⁸². At the hospital-level, facilities with higher percentage of Black patients are found to be under considerable financial strain, leading to poor quality and safety outcomes⁸¹.

Payor:

Clinical outcomes are known to vary by the type of insurer financing the particular episode of care⁸³⁻⁸⁶. The choice of a distinct set of hospitals by uninsured and Medicaid patients and a disproportionate volume of disadvantaged patients with certain physicians are some factors that influence this variation. Moreover, studies have also shown that within hospitals the quality of care differs by the insurance type⁸⁷. This may be partly attributed to the differences in reimbursement by insurers, differences in physicians who serve patients of a certain insurance type, and lower utilization of latest modalities of care for Medicaid patients because of reimbursement constraints.

Admission Type:

The type of admission – elective, urgent or emergent, is a determinant of outcomes and quality of care^{88, 89}. Patients admitted to hospitals on urgent or emergent basis are known to be at a greater risk of developing complications⁹⁰. The severity of illness,

intensity of treatment protocols, and greater resource use are some of the factors that explain this association.

Admission Source:

The source of admission, such as home, another hospital, and emergency department are predictors for complications during both medical and surgical hospitalizations⁹¹⁻⁹³. Patients who were previously admitted to other healthcare facilities may have undergone significant medical and surgical management, thereby providing opportunity for care-related harm. Furthermore, the severity of illness may have aggravated in these patients, which may have warranted their transfer to another health care facility. Thus, given their compromised physiologic status, such patients are more likely to be at risk of complications as compared to patients who are electively admitted from home.

Comorbidities:

Presence of comorbidities at the time of admission is an important predictor of clinical outcomes and complications, especially in older patients⁹⁴⁻⁹⁶. Decreased physiological reserve, decreased cognition, need for large number of medications and procedures, and involvement of several clinicians in the care process are some of the factors that contribute to the vulnerability of older patients to medical errors.

Surgical procedure:

Patients undergoing surgeries are at a greater risk of death, surgical complications such as surgical site infections, and medical complications such as myocardial infarction^{97, 98}.

Controlling for the occurrence of a major, minor or no surgical procedure will partially explain the variation in HACs.

Weekend admission:

Patients admitted to hospitals on a weekend tend to have poorer outcomes as compared to those admitted on weekdays in certain instances ⁹⁹⁻¹⁰¹. Limited patient access and staffing ratios are some of the factors that explain 'weekend effects'.

Median household income in zip code:

Patients living in a particular location are likely to have similar health status, health risks, and availability of healthcare providers. Thus, patients living in neighborhoods with lower household incomes are more likely to have poorer health outcomes, especially when hospitalized ¹⁰². Thus, the median household income variable is a proxy for the patient's health status and the quality of accessible care.

Hospital-level factors

Hospital size and occupancy:

Hospitals with larger number of beds and higher occupancy rates were found to have lower adjusted mortality rates and better care quality ^{68, 69, 103}. Higher occupancy rates may be a proxy for good financial health of the hospital, which in turn may translate into larger investments in QI initiatives. Furthermore, with higher volumes, the 'practice makes perfect' principle may result in superior outcomes ⁶⁷.

Urban or rural setting:

Rural hospitals have often been found to deliver lower quality care^{68, 104} as compared to hospitals located in urban areas. Inadequate work force and material resources have often explained the quality in rural hospitals. In contrast, rural hospitals may also transfer severely ill patients to tertiary-care facilities, and this may result in rural hospitals having a larger proportion of less severely ill patients as compared to urban hospitals, leading to better performance on quality metrics⁶⁷.

Ownership:

Studies have documented variation in the quality of care between for-profit (FP) and not-for-profit (NFP) hospitals, with NFP hospitals performing better on several metrics¹⁰⁵⁻¹⁰⁷. Several factors explain the superior performance of NFP, including the role of NFP as quality-quantity maximizers, their altruistic philosophy. In contrast, the lack of adequate finances after diversion of profits to stakeholders, and location in areas with low quality are some of the mechanisms explaining the poor performance of FPs.

Financial health:

Several studies have linked the financial stability of the hospital to better quality and safety outcomes^{69, 108}. Studies have demonstrated that patients admitted to hospitals with inadequate resources are more likely to receive poor quality care¹⁰⁹⁻¹¹¹. Adequate financial resources empowers the hospital to make investments in QI initiatives, attracting and retaining qualified workforce, providing tools and resources such as IT to enable the staff to perform their duties adequately.

Teaching status:

The teaching status of a hospital is strongly linked to the quality of care, with teaching hospitals performing better on quality and safety metrics as compared to non-teaching hospitals^{68, 112, 113}. This can be partly explained by factors such as higher qualifications and experience of the workforce, availability of superior tools and technology and urban location. In contrast, a larger volume of acutely ill patients may also result in higher incidence of iatrogenic injury among patients, thereby resulting in poor quality care⁶⁷.

Patient-payer mix:

Hospitals that care for disproportionately high number of disadvantaged patients often perform poorly on certain quality metrics. The likelihood of HACs among safety-net hospitals is elaborately discussed in the third Specific Aim (see Section 1.6.4).

Electronic Health Record capabilities:

The adoption and use of EHRs is an integral part of QI initiatives. The role of EHRs in reducing the likelihood of HACs is the subject of the Specific Aim 2.1 and is discussed in detail in Section 1.6.4.

Market competition:

Market competition has varying associations with the quality of care delivered. Studies have demonstrated improvement in quality and reduction in costs with increased market competition^{114, 115}. In contrast, studies have demonstrated an association between increasing competition and declining performance on quality metrics^{116, 117}. Improvement in efficiency and resources can be linked to improvement in outcomes with increased competition, whereas lack of quality directives and limited financial margins by the

payers for patients being sought after are some factors that may explain the adverse impact.

1.6.2. Specific Aim 2 – Hypothesis 2.1

Impact of P4P policies on quality in hospitals

P4P has been identified as the ‘single most promising approach’ to improve the quality of care while controlling costs ^{118, 119}. The rationale for P4P programs is to provide hospitals with incentives that will motivate them to bring about changes in their structures and processes, and that are likely to result in improved outcomes ^{35, 120}.

Despite the intuitive appeal, face validity, and theoretical soundness of P4P programs, the response to these programs in the US has been mixed ^{34, 121, 122}. The largest P4P program in recent times, the Medicare Premier Hospital Quality Incentive Demonstration (HQID), evaluated the impact of bonuses on 5 clinical conditions, namely acute myocardial infarction (AMI), heart failure, community-acquired pneumonia, coronary artery bypass graft, and hip and knee replacement. Short-term baseline analysis demonstrated modest improvement in composite process measures, for hospitals involved in both P4P and public reporting, as compared to hospitals that participated only in public reporting ¹²³. Improvements were largest for hospitals that received larger bonuses, had a better financial standing and were located in less competitive markets ¹²⁴. However, an in-depth analysis of individual AMI measures did not provide evidence for an improvement in performance associated with the HQID ¹²⁵. There is evidence of the program reducing mortality and growth in costs, and a small influence has been demonstrated on the outlier classification ¹²⁶. Long-term analysis of the HQID found minimal effects on compliance to process measures and no evidence of favorable influence on outcome measures ^{35, 124, 125, 127}. In contrast, Sutton et al. found that the 2008 Advancing Quality P4P program in the United Kingdom, which was based on the

US HQID program, resulted in a significant absolute decline in mortality by 1.3 percentage points and a relative decline of 6% for the duration of the program ¹²⁸. This decline was particularly pronounced for pneumonia-related mortality which significantly declined by 1.9 percentage points. Long-term analysis of this initiative did not identify sustained significant decline in mortality. This is partly explained by positive, quality-related spillover effects on the comparator group in anticipation of impending quality legislations, and on conditions that were used as control groups ¹²⁹.

A number of factors explain the variation in the response of hospitals to the P4P programs. Significant among these are, the magnitude of the incentives at stake, the structure of the incentives, the choice of appropriate metrics, and other contextual factors such as healthcare system, payer, provider and patient characteristics ^{35, 122, 124, 130-133}.

While rewarding hospitals and clinicians with bonuses for meeting certain standards of care or demonstrating a predetermined improvement in performance are popular versions of P4P ('carrot'), the approach of reducing reimbursements for poor quality care ('stick') has shown promise. Not paying for non-performance or poor quality care is considered the mirror image of P4P ³⁶. In this approach, penalties are levied when hospital performance falls below a certain threshold or hospitals are not reimbursed for the care of conditions that were not present on admission ^{7, 134}. HealthPartners, a not-for-profit health maintenance organization (HMO) based in Minnesota, was among the first health care purchasers to adopt this approach. In January 2005, HealthPartners reduced payments for 'Never Events' that developed during a patient's stay in the hospital ¹³⁵.

This model of controlling adverse events was soon adopted by several public payers including Minnesota's insurance programs, other states, and the federal government ³³.

Public programs that introduced penalties include the HQID program and the Hospital Readmission Reduction program. The HQID program introduced penalties for non-performing hospitals in Year 3 of the program ¹²³. In FY 2006, hospitals whose performance scores fell in the bottom 20% of the first-year scores were levied penalties ranging for 1 to 2% of their base Medicare payment. The HRR program imposes penalties between 1 to 3% of the base Medicare rates to hospitals based on their readmission adjustment factor for both Medicare and non-Medicare patients ¹³⁶.

Impact of HAC Program on HAC rates in hospitals

I have identified ten quantitative and six qualitative studies that evaluate the impact of the HAC Program (see Table 2).

Table 2: Studies evaluating the impact of the HAC Program on HAC rates

Author	Study Period	Post-Program	Scope of HACs	Scope	Data
Quantitative Studies					
McNair	2005 to 2006	No	All	National and California	Administrative: HCUP, California discharge data
McNutt	2005 to 2007	No	All	National	Administrative: University HealthSystem Consortium
McHugh	2006	No	All	National	Administrative: Medicare Claims
Stone	2008 and 2010	Yes	HAIs (excluding SSIs)	California	Hospital rates
Lee	2006 to 2011	Yes	HAIs	National	Administrative: Nationwide Inpatient Sample
Meddings	2007 and 2009	Yes	CAUTIs	Michigan	Administrative: HCUP Michigan
Gidwani	2007 to 2009	Yes	DVT / PE	National	Administrative: Nationwide Inpatient Sample
Sculler	2005 to 2009	Yes	UTI	National	Administrative: Nationwide Inpatient Sample
Waters	2006 to 2010	Yes	CAUTI, CLABSI, Falls and Pressure Ulcers	National	Clinical: National Database for Nursing Quality Indicators
Vaz	2007 to 2013	Yes	CLABSI	National	Clinical: National Healthcare Safety Network
Qualitative Studies					

Author	Study Period	Post-Program	Scope of HACs	Scope	Data
Hoff	2009 to 2010	Yes	HAIs	National	Survey of Infection Preventionists (IPs)
Krein	2005 and 2009	Yes	HAIs	National	Survey of IPs
Stone	2008 and 2010	Yes	HAIs (excluding SSIs)	California	Survey of IPs
Hartmann	2009 to 2010	Yes	HAIs	National	Survey of IPs
Lee	2010	Yes	HAIs	National	Survey of IPs
McHugh	2009	Yes	All	National	Survey of Safety-Net Hospital Administrators

Abbreviations: HCUP: Healthcare Cost and Utilization Project; HAI: Healthcare-associated infection; SSI: Surgical site infection; CAUTI: Catheter-associated urinary tract infection; DVT/PE: Deep vein thrombosis / pulmonary embolism; CLABSI: Catheter-associated bloodstream infection

McNair et al used 2006 Patient Discharge Dataset of the California Office of State Health Planning and Development to simulate the potential impact of the HAC Program in California and the 2005 HCUP Nationwide Inpatient Sample to compare the prevalence of HACs in California with national rates ¹³⁷. The study found that 0.11% of the 767,995 Medicare discharges in California had HAC-related codes and an estimated \$92,000 to \$227,000 would be saved by not reimbursing Californian hospitals for the care of HACs. Nationally, annual savings were estimated to be between \$1.1 to \$2.7 million, and consequently the Program was expected to have limited impact.

McNutt et al analyzed University HealthSystem Consortium data from 86 academic medical centers for the duration between 2005 to 2007 ¹³⁸. Using bootstrap methods to estimate the proportion of HACs, the authors found that the HAC Program may reassign MS-DRGs for 7.5% of cases with HAC-associated secondary diagnoses codes, and non-payment for these reassignments is expected to annually save \$25 million for the CMS. The difference in the financial impacts estimated by McNutt et al. and McNair et al. primarily result from the scope of cohort. The study by McNutt et al. includes all discharges regardless of their payer whereas the study by McNair et al. includes only the discharges reimbursed by Medicare.

McHugh et al used Medicare claims and AHA survey database for 2006 to study the difference in HAC rates between safety-net and non-safety-net hospitals ¹³⁹. Using t-tests and ANOVA, they found significantly higher HAC rates among hospitals that served uninsured, Medicaid and other vulnerable patients. Additionally, the study identified an inverse relationship between the total financial margin of the hospital and HAC rates.

HAC rates were also found to be higher among urban, public, large (>300 beds) and teaching hospitals. The higher HAC rates in safety-net hospitals are likely to strain their financial status, with little opportunity to cross-subsidize. Given these findings, the authors identified structural (such as nurse staffing and culture), and patient-related factors as key in preventing HACs from occurring.

Stone et al linked incidence rates of HAIs in the ICUs of 200 Californian hospitals in 2008 and 2010 with qualitative data gathered from semi-structured interviews with infection preventionists of these hospitals to assess the impact of the HAC Program on the structure, processes, and outcomes relevant to the HAIs that are included in the Program ¹⁴⁰. Survey results found the infection preventionists spent more time on surveillance activities following the implementation of the Program, resulting in decreased time for educational activities. The study also identified significant improvements in adherence to EBPs for CLABSI and CAUTI, and a decline in the rates for CLABI between the two periods.

Lee et al studied the impact of the HAC Program on infections in US hospitals using clinical data from CDC's National Healthcare Safety Network (NHSN) for the years 2006 to 2011 ³⁴. They used an interrupted time series design along with comparison series for studying the trends in catheter-associated blood stream infections and urinary tract infections, after controlling for trends in ventilator-associated pneumonia. The researchers did not find evidence of the HAC Program having a measurable impact on the metrics of interest, after controlling for relevant variables.

Meddings et al studied discharges from Michigan hospitals that took place in 2007 and 2009 to assess the impact of the HAC Program on rates and reimbursements for CAUTIs ¹⁴¹. They compared these rates with rates of non-catheter associated UTIs for the same period. Using t-tests, the researchers found negligible change in the rates of CAUTIs with the implementation of the HAC Program. The low rates of CAUTIs found in the study (mean of 0.09% in 2007 and mean of 0.14% in 2009) was largely explained by the uncommon use of the ICD-9-CM code 996.64, which is required for the identification of an UTI as a CAUTI per the Program. This was mainly due to the recording of the catheter-related information in nursing notes, whereas the information used for HAC coding was derived from physician notes. Furthermore, significant differences in the coding guidelines for billing as compared to those used for medical records and registries further augmented the number of false-negative cases. Given the complex coding requirements for identification of CAUTIs, the study estimated an annual loss of only 0.06% of annual Medicare payments that could be attributed to CAUTIs.

Schuller et al ¹⁴² used the Nationwide Inpatient Sample to study the risk of CAUTIs from 2005 to 2009. They did not identify significant changes in the risk of CAUTIs with the implementation of the Program. They suggest that the low financial implications of the Program are a key factor in the non-response of hospitals to the Program. They make several suggestions including adherence to protocols and evidence-based measures, focusing on physicians and other members of the clinical team, and carrying out cost-benefit analyses as potential strategies to make hospitals sensitive to the Program.

Gidwani et al ¹⁴³ used the 2007-2009 Nationwide Inpatient Sample to assess the impact of the Program on the risk of deep vein thrombosis or pulmonary embolism (DVT/PE) in patients undergoing hip and knee replacements. They identified significant favorable effects of the Program, and attributed it to consistent use of prophylaxis for patients undergoing these surgeries. Their findings include positive spillover effects of the Program to elderly inpatient stays not financed by Medicare.

Waters et al ¹⁴⁴ used the National Database for Nursing Quality Indicators from 2006 to 2010 to identify unit-level effects of the Program. They identified significant Program effects on the CAUTI and CLABSI rates. These effects were attributed to the availability of evidence-based guidelines, standardization of practices, and relative preventability of these conditions. The study did not identify significant effects on rates of pressure ulcers and falls.

Vaz et al ¹⁴⁵ used 2007-2013 NHSN data to study the impact of the Program on CLABSI in safety-net and non-safety-net hospitals. They did not identify significant changes in the rates of CLABSIs. These findings were attributed to the limited financial impact of the Program, a generalized downward trend in CLABSIs because of quality awareness, and increased efforts by safety-net hospitals to meet quality standards.

Hartmann et al sampled 36 hospitals from the AHA database and conducted semi-structured telephone interviews with infection preventionists from these hospitals between 2009-2010 ¹³⁰. Questions ranged from HAI-related practices prevalent in their hospitals to the response of their hospitals to the HAC Program and its consequences.

The three main themes that evolved from the responses were change in systems, diversion of resources from other important activities and change in staff activities relating to surveillance.

Hoff et al used the aforementioned survey to examine the role of leadership and culture in influencing the hospital response to the Program ¹⁴⁶. They found that top leadership and clinical staff involvement, and a proactive organization culture were key in optimizing the effectiveness of the Program. The importance accorded to the HAC Program by the key stakeholders, the role of the Program in enhancing the educational efforts and the alignment of the Program with other QI efforts prevailing in the hospital significantly contributed in enhancing organizational awareness towards QI in general and HAIs in particular.

Krein et al conducted longitudinal surveys with infection preventionists from non-federal and VA (used as comparator) hospitals in 2005 and 2009 to examine (i) the change in the adoption of EBPs for preventing HAIs such as CLABSIs, VAPs, and CAUTIs, and (ii) the change in the priority assigned to the prevention of these HAIs ¹⁴⁷. The study found that the HAC Program was associated with an increase in priority accorded to the prevention practices in non-federal hospitals as compared to VA hospitals. However, this increase in priority did not translate into an increase in the adoption of EBPs and considerable variations in practices persisted.

Lee et al surveyed infections preventionists from a sample of 500 acute care hospitals in the US to determine the perceived impact of the HAC Program on the steps taken and

resources available to reduce HAIs and understand the unintended consequences of the Program ¹⁴⁸. The study results were largely promising and found that the HAC Program had increased the attention of hospital stakeholders towards HAIs included in the Program, and improved relevant practice patterns. However, few hospitals also acknowledged the diversion of focus away from infections not included in the Program, and the adoption of 'defensive' measures to prevent HAIs from occurring.

McHugh et al telephonically surveyed 120 Chief Financial Officers and Chief Safety Officers of randomly sampled safety-net hospitals to understand the impact of the HAC Program in the first year of its implementation ¹⁴⁹. They found that the Program had caught the attention of hospital leaders and governing boards, and provided added motivation to prevent HACs from occurring. However, given the limited financial impact of the Program, few hospitals had brought in elaborate QI measures to aid prevention, and several hospitals focused their resources towards coding and documentation practices to prevent penalties. Furthermore, the impending expansion of the Program to include Medicaid patients, and the focus on HACs through other QI programs such as the Joint Commission accreditation may have been responsible for the heightened awareness towards the Program.

1.6.3. Specific Aim 2 – Hypothesis 2.2

Specific Aim 2.2 evaluates the impact of the Program across hospitals with varying financial strength. This section summarizes literature that motivates this Aim. I begin by synthesizing literature on the relation between the financial strength of a hospital and the quality of care it renders; followed by a review of the impact of P4P programs among hospitals with varying financial strength; and finally literature on the specific impact of the HAC Program among these hospitals.

Association of hospital financial performance with quality and safety of care

The provision of high-quality care needs considerable financial investments. These monetary resources are essential to adopt, maintain and update QI initiatives such as recruitment of qualified and skilled staff in adequate numbers, procurement of information technology and the adoption and integration of evidence-based processes within routine workflows. Studies have demonstrated that hospitals in poor financial health tend to control their expenses by limiting investments in QI. Consequently, patients who are admitted to such hospitals often receive poor quality care ^{109-111, 150}.

Impact of P4P policies in SN hospitals

Recent studies suggest that the care for racial and socio-economically disadvantaged patients is concentrated in a small proportion of hospitals, commonly known as Safety-Net (SN) hospitals, several of which perform poorly on quality metrics ¹⁵¹⁻¹⁵³. SN hospitals “organize and deliver a significant level of health care and other health-related services to uninsured, Medicaid, and other vulnerable patients” ¹⁵⁴.

SN hospitals are known to often deliver lower quality of care as compared to non-SN hospitals^{134, 151, 155, 156}. While P4P programs that align bonuses or penalties to hospital performance are expected to improve the quality of care^{123, 156-160}, there are concerns that P4P also has a paradoxical potential to aggravate existing disparities^{134, 157, 161-167} between SN and non-SN hospitals. By widening the gaps between hospitals that have a disproportionately high load of underserved patients, P4P programs may effectively deny SN hospitals the economic benefits of P4P¹⁵⁵.

Werner et al¹³⁴ simulated the impact of CMS' HQID incentives on the care of acute myocardial infarction, heart failure, and pneumonia in national hospitals with varying percentages of Medicaid patients. They found that hospitals with higher percentages of Medicaid patients made smaller improvements in their performance scores as compared to hospitals with lower percentage of Medicaid patients. Thus, these hospitals were less likely to receive financial bonuses from the program, further limiting the resources available for QI initiatives.

In contrast, Jha et al¹⁵⁶ limited their study to the sample of hospitals that participated in the HQID program and used non-participating hospitals as a comparison group. They found that financial incentives of P4P programs were useful in stimulating QI in hospitals. Hospitals that had a larger volume of poor patients, considerably improved their quality performance in response to the incentives of the HQID program, and were at par with hospitals that had a few poor patients by the end of the first phase of the program.

Ryan ¹⁶⁸ examined the association between the proportions of disadvantaged patients in a hospital and the incentive payments from the first year of the Hospital VBP Program. He used the Disproportionate Share Hospital (DSH) index to represent the magnitude of socioeconomically disadvantaged patients in a hospital, and found that hospitals with higher DSH index performed worse in comparison to those with lower DSH index. While the small size of payments was unlikely to aggravate disparities in the first year, it is expected that the combined effect of HVBP and the Hospital Readmission Reduction Program is likely adversely influence the quality of care in SN hospitals.

A number of mechanisms by which the unintended negative consequences of P4P take root in SN hospitals have been identified. The financial status of the hospital and the design of P4P programs are important factor that place SN hospitals at an increased risk of disparities ^{149, 154, 157, 160, 165, 169-173}. Other mechanisms such as a 'one size fits all' approach of P4P programs without customization to needs of the underserved population, 'teaching to the test' approach by focusing only on metrics that are tied to payments while ignoring those that are associated with disparities, inadequate risk-adjustment of outcomes, and 'cherry picking' by avoiding patients with adverse health profiles who may be threats to the quality score of a hospital are some of the several mechanisms by which P4P may perpetuate disparities in the quality of care ^{133, 157, 161, 162, 169, 171, 173}.

Impact of HAC Program in Safety-Net Hospitals

There is little empirical evidence on the differential impact of the HAC Program in SN and non-SN hospitals. Two relevant studies ^{145, 149} are discussed in Section 1.6.2.

1.6.4. Specific Aim 2 – Hypothesis 2.3

Impact of EHRs on quality and safety

An extensive body of literature supports the role of EHRs in improving quality, enhancing safety, and reducing the costs of care delivered in ¹⁷⁴⁻¹⁸³. In a recent review, Buntin et al. ¹⁸⁴ found that 92% of studies reviewed had positive findings about the impact of health IT. This review included studies that found improvements in safety-related activities such as medication prescribing ¹⁸⁵⁻¹⁸⁸, documentation and communication ¹⁸⁹⁻¹⁹², and HAI surveillance with the use of EHRs ¹⁹³⁻¹⁹⁶.

Studies specific to safety conditions included in the HAC Program have shown promising results. Studies have demonstrated a significant improvement in completeness and accuracy of pressure ulcer documentation with the implementation of EHRs ¹⁹⁷⁻¹⁹⁹. Implementing EHRs has also been associated with a decline in the incidence ²⁰⁰ and prevalence of hospital-acquired pressure ulcers ²⁰¹, and in the incidence of falls ^{202, 203}. EHRs have also been associated with increased adherence to recommended diabetes management guidelines ²⁰⁴⁻²⁰⁸ and improvement in blood glucose levels ^{208, 209}. However, some studies did not find evidence of improvement in blood glucose levels with the use of EHRs ^{206, 210, 211}. In this regard, better integration of the technology with routine workflows, and enhanced patient education and activation for utilizing EHRs were identified as areas of improvement. Several studies have demonstrated the utility of EHR-enabled algorithms to support HAI surveillance ²¹²⁻²¹⁴. For example, the use of algorithms to identify CAUTIs from cultures reduced the need for manual surveillance by 97% ²¹⁵. Welden et al. used EHRs to provide clinical decision support and evidence-based care resulting in decreased utilization of catheters and reduced catheter-usage

duration ²¹⁶. EHR functionalities have favorably influenced the identification of patients at risk of developing VTE, compliance to VTE prophylaxis protocols, and, consequently, decreased incidence of VTE ²¹⁷⁻²²⁴.

Impact of P4P policies in hospitals with EHRs

The competitive environment that is introduced by P4P policies such as the HAC Program is expected to motivate hospitals to maximize the use of IT resources at their disposal. However, there is lack of empirical evidence that addresses the interaction between prevailing P4P policies and contextual factors such as use of EHRs in hospitals ¹⁸³. The study by Jha et al. is among the few studies that evaluated the impact of EHR adoption and use on quality metrics among safety-net hospitals ²²⁵. Their findings suggest that EHR adoption motivated significant improvements in quality in safety-net hospitals. Furthermore, small, nonteaching, and rural hospitals were found to be slow adopters of EHRs ²²⁶. Considering that these groups of hospitals are known to be less effective in their response to P4P policies, the findings of these studies support the expectation that the response to such P4P policies may be modified by the capabilities and extent of use of EHRs.

1.6.5. Gaps in HAC Program literature

I have identified sixteen empirical studies (ten quantitative and six qualitative) that investigate the impact of the HAC Program on HAC rates. This dissertation will address the drawbacks of current quantitative studies as follows:

Longitudinal study duration

Several important quality reforms, especially those resulting from the Affordable Care Act, came into effect at the start of the current decade. It is important that these reforms are accounted for in the analysis to identify the independent effects of the HAC Program. With the exception of the studies by Lee et al and Vaz et al ^{34, 145}, most of the quantitative studies have used data only from the previous decade. This dissertation uses data from 2005 to 2012, thereby accounting for secular trends in outcomes and contemporaneous reforms that may confound the impact of the HAC Program.

Present-on-Admission indicator

Of the ten quantitative studies that evaluated the impact of HAC Program on HAC rates, three studies simulated the impact of the Program using data from before the HAC Program was implemented ¹³⁷⁻¹³⁹. Thus, their analyses are limited by the fact that POA indicators were not included in the datasets (these indicators were mandated for discharges after October 1, 2007), and hence the studies simulated the probabilities as to which conditions may have been acquired following admission of the patient to a hospital.

Furthermore, two other studies have used the Nationwide Inpatient Sample to evaluate the Program ^{142, 143}. However, the POA indicators are not included in this dataset, thereby limiting the validity of their findings. This dissertation addresses these gaps by using the New York (NY) State Inpatient Databases for New York, which not only report the POA indicator for each diagnosis, but are one of only two databases (the other one being the California State Inpatient Database) that have reported the POA indicator since the mid-90s.

Outcomes studied

Hospital-acquired infections have been the target of several federal-, state-, and local-level reforms, and most the HAC Program-related studies have focused on evaluating the impact of the Program on these conditions. Only two studies ^{143, 144} have evaluated the impact of the Program on non-infection outcomes such as DVT/PE, falls, and pressure ulcers. This dissertation addresses this gap by evaluating the impact of the Program on all HACs that are within the scope of the Program, and demonstrate reasonable variation.

Types of data

Only two studies have used clinical data to evaluate the impact of the Program ^{34, 144}. These studies utilized clinical data from the Preventing Infectious Complications by Adjusting Payment Project ²²⁷, and the National Database of Nursing Quality Indicators ²²⁸. The key advantages of using these clinical datasets are the sensitivity and specificity of the outcomes ^{229, 230} abstracted from clinical records. However, the use of these datasets also has certain drawbacks. First, these datasets contain data from hospitals

that voluntarily participate in these programs. These participating hospitals are likely to be quality oriented, and to have the workforce and material resources to participate in such Programs. This non-randomized nature of the study cohort is likely to bias the study findings. Second, these databases contain unit-level (such as specialty wards or intensive care units) data, and thus do not contain details on patient-level factors. Thus, by not adjusting for patient-level risk factors, the studies may have missed important Program-associated changes in the patient-level HAC variance. Importantly, the aggregate nature of the data does not support the identification of the insurance status of the patient. It is likely that the aggregate units may have varied proportions of Medicare patients, which cannot be accounted for in the analysis. This dissertation addresses these drawbacks of previous studies by using the HCUP database that includes all inpatient Medicare observations from New York State thereby limiting the selection bias, and that contains patient-level data to adjust for patient risk factors. Importantly, even though the HCUP database is an administrative database and has limitations of sensitivity and specificity as compared to clinical data, the CMS uses similar administrative datasets to assess the financial disincentives for hospitals.

Study design

Only three studies use a control group to evaluate the Program effect ^{34, 143, 145}. Importantly, the control groups that are used in these studies are likely to be subject to the spillover effects from the Program thereby limiting the identification of the independent effects of the Program. Furthermore, these control groups do not model the financial risk that a hospital is likely to perceive from the Program. This dissertation addresses this concern by interchangeably using the lower MUR quartiles as control

groups to model the financial risk from the Program (see Section 1.5.3.), and difference-in-difference (DID) estimation to control for secular trends

Differential Program effects

The financial status of a hospital and its digital capabilities are key factors in influencing the hospital response to P4P Programs and the quality of care rendered. This dissertation evaluates the differential effect of the Program across hospitals with varying financial strength and EHR capabilities – thus contributing to a limited knowledge base on the Program.

Chapter 2 - METHODOLOGY

2.1. Data sources

I have used a combination of individual-, and hospital-level databases to address the research questions.

2.1.1. Healthcare Cost and Utilization Project (HCUP) databases for NY

The Healthcare Cost and Utilization Project's (HCUP) State Inpatient Databases (SID) for NY contain the universe of discharge records for patients discharged from NY hospitals ^{231, 232}. NY Department of Health provides the data for HCUP NY SID. The SID includes patient-level information derived from billing records for insured (public and private) and uninsured patients. These are organized in five different files, namely the core file, charges file, AHA linkage file, Diagnosis and Procedure Groups file, and the Disease Severity Measures file. Quality of the data is maintained by conducting internal consistency and edit checks. Importantly, since the mid-1990s, the NY SID includes the Present on Admission (POA) indicator that is used for identifying patient safety events that were acquired after admission. In NY, 83% of discharges passed checks that examined the face validity of the POA indicators ²³³. Moreover, continual efforts are in place to communicate the value of POA indicators to hospitals and coders ²³⁴. The SIDs have been extensively used to evaluate various aspects of the cost, quality, and delivery of care in the US health system. Since these datasets are derived from billing data, a key limitation of these datasets is the absence of clinical information that enhances the sensitivity and specificity of the findings.

2.1.2. CMS Impact files

CMS Impact files provide estimates of the impact of various Program changes introduced in the IPPS through the proposed and final rules published in the Federal Register ²³⁵. These files contain variables required for the computing of the reimbursements for hospitals (operating and capital base payment rates) under the IPPS. The files include variables such as geographic factors, wage index, DRG weights, and disproportionate share patients.

2.1.3. CMS Provider of Services files

CMS' Provider of Services ²³⁶ files contain data on facility characteristics, services provided, human resource composition and several other structural indicators for hospitals and other healthcare institutions.

2.1.4. Dartmouth Atlas databases

The focus of the Dartmouth Atlas Project has been to study variations in the availability and utilization of resources across the country ²³⁷. The Dartmouth Atlas geographic crosswalk files were used to assign NY hospitals to their respective Hospital Referral Regions (HRR) and Hospital Service Areas (HSA).

2.1.5. Hospital Information and Management Systems Society (HIMSS) databases

HIMSS and HIMSS Analytics LLC jointly maintain the HIMSS databases ²³⁸. These databases contain variables on hospital characteristics and health IT products, processes and use in more than 4,000 US hospitals. The database was used to identify the preparedness of the hospital to fulfill CMS' 'Meaningful Use' criteria.

2.1.6. CMS EHR Incentive Program files

The EHR Incentive Program for the adoption and Meaningful Use of EHRs was mandated by the HITECH Act of 2009 ²³⁹. The Program files contain details of the fulfillment and payments made to eligible hospitals and providers that have attested to the Meaningful Use criteria for various stages of the Program.

2.1.7. CMS Cost Reports

CMS's Cost Reports ²⁴⁰ for hospitals contain details of hospital costs and charges, utilization, characteristics, Medicare settlement data and other financial details.

2.1.8. AHA ID and CMS Provider ID linkage files

Hospital names, geographic locations and AHA's online hospital lookup ²⁴¹ were used to create a crosswalk file that links Medicare Provider IDs with AHA IDs for hospitals in NY. This enabled the linkage of hospital-level files that contain Medicare IDs for hospitals with the patient-level HCUP files that contain AHA IDs.

2.2. Cohort specification

A diagrammatic representation of the algorithm used to identify the final cohort is presented in Figure 4.

. The following criteria were used to define the cohort:

- The cohort was limited to discharges ≥ 65 years of age and ≤ 99 years of age, and to discharges for which Medicare Part A was the primary, secondary or tertiary payer.
- In order to account for the heterogeneity among the observations, the dataset was divided into medical and surgical cohorts of interest. For the medical cohort, discharges with acute myocardial infarction, pneumonia, congestive heart failure²⁴² and stroke^{243, 244} as the primary diagnosis were identified using ICD-9-CM diagnosis codes. For the surgical cohort, discharges with ICD-9 procedure codes that are specified by the HAC Program were retained. The procedures of interest are CABG, spine surgeries, and hip and knee replacement. Table 3 provides the details of the ICD-9-CM diagnosis and procedure codes used for creating the cohort.
- Stays with discharge dates prior to October 1, 2008 were classified as the pre-Program cohort. Discharges from 2008 that were missing data on the discharge month were excluded from the cohort. Stays with discharge dates from October 1, 2008 to December 2012 were included in the post-Program cohort.
- The resulting patient-level dataset was merged with the hospital-level datasets using the AHA ID to retain observations from hospitals reimbursed by the IPPS and for whom linking identifiers existed in the two databases.

Figure 4: Cohort identification algorithm

Table 3: ICD-9-CM Diagnosis and Procedure codes used for cohort identification

Cohort	HAC	ICD-9-CM Diagnosis and Procedure Codes
Medical	Acute myocardial infarction	410.00, 410.01, 410.10, 410.11, 410.20, 410.21, 410.30, 410.31, 410.40, 410.41, 410.50, 410.51, 410.60, 410.61, 410.70, 410.71, 410.80, 410.81, 410.90, 410.91
Medical	Congestive heart failure	402.01, 402.11, 402.91, 404.01, 404.03, 404.11, 404.13, 404.91, 428.0, 428.1, 428.20, 428.21, 428.22, 428.23, 428.30, 428.31, 428.32, 428.33, 428.40, 428.41, 428.42, 428.43, 428.9
Medical	Pneumonia	480.0, 480.1, 480.2, 480.3, 480.8, 480.9, 481, 482.0, 482.1, 482.2, 482.30, 482.31, 482.32, 482.39, 482.40, 482.41, 482.42, 482.49, 482.81, 482.82, 482.83, 482.84, 482.89, 482.9, 483.0, 483.1, 483.8, 485, 486, 487.0, 488.11
Medical	Stroke	430, 431, 432.0, 432.1, 432.9, 433.00, 433.01, 433.10, 433.11, 433.20, 433.21, 433.30, 433.31, 433.80, 433.81, 433.90, 433.91, 434.00, 434.01, 434.10, 434.11, 434.90, 434.91, 436, 438.0, 438.10, 438.11, 438.12, 438.13, 438.14, 438.19, 438.20, 438.21, 438.22, 438.30, 438.31, 438.32, 438.40, 438.41, 438.42, 438.50, 438.51, 438.52, 438.53, 438.6, 438.7, 438.81, 438.82, 438.83, 438.84, 438.85, 438.89, 438.9
Surgical	Coronary Artery Bypass Graft (CABG)	36.10-36.19
Surgical	Certain orthopedic procedures (Spine, neck, shoulder, elbow)	81.01-81.08, 81.23-81.24, 81.31-81.38, 81.83 or 81.85
Surgical	Bariatric surgery for obesity (Laparoscopic gastric bypass, gastroenterostomy, laparoscopic gastric restrictive surgery)	44.38, 44.39, 44.95
Surgical	Certain orthopedic procedures (Total knee replacement, hip replacement)	00.85-00.87, 81.51-81.52, or 81.54

Abbreviations: ICD-9-CM: *International Classification of Diseases, Ninth Revision, Clinical Modification*

2.3. Datasets and variables

2.3.1. Dataset and variable mapping

Table 4 presents a summary of the datasets, variables, and variable specification used in the analyses.

Table 4: Dataset and variable linkages

Variable Category	Variable	Data Source	Level	Hospital Linkage	Specification
Dependent	HACs	HCUP SID	Patient	AHA ID	<ul style="list-style-type: none"> - Categorized as binary indicator (see Section 2.3.2 for details) - Primary outcome: Any-or-none indicator for each cohort - Secondary outcomes: Each of the 6 HACs for each cohort, surgical site infections in the CABG and spine sub-cohorts, deep vein thrombosis/pulmonary embolism in the hip and knee cohorts
Key independent	Program phase	HCUP SID	Patient	AHA ID	- Categorized as a binary indicator (see Section 2.3.3 for details)
	Medicare Utilization Ratio	Impact files	Hospital	Medicare Provider ID	- Categorized into hospital-level quartiles (see Section 2.3.3 for details)
	Operating profit margin	Cost report			- Categorized as positive or negative operating margin (see Section 2.3.4 for details)
	DPP	Impact files			- Categorized as above or below median DPP (see Section 2.3.4 for details)
	EMRAM Score	HIMSS			- Categorized from Stage 0 to Stage 3 and above (see Section 2.3.4 for details)
Covariates: Patient	Age	HCUP SID	Patient	AHA ID	- Categorized as <70, >=70 and <80, >=80 and <90, >=90
	Sex				- Categorized as male or female

Variable Category	Variable	Data Source	Level	Hospital Linkage	Specification
	Race and ethnicity				- Categorized as White non-Hispanic, Black non-Hispanic, Hispanic, Other non-Hispanic
	Admission type				- Categorized as emergent/urgent/trauma, elective
	Primary payer				- Categorized as Medicare, private, other
	Admission source				- Categorized as transfer from a healthcare facility, admitted directly from home/doctor's office
	Surgical procedure				- Categorized as none, major, minor
	Weekend admission				- Categorized as yes, no
	Median income quartile for zip code				- Categorized as quartile 1 to 4
	Comorbidities from Elixhauser algorithm				- 29 binary indicators for each comorbidity
	Discharge year				- Categorized as 2005 to 2012
	Cohort and sub-cohort of interest				- Categorized based on the primary diagnosis of interest for the medical cohort, and the surgeries of interest for the surgical cohort (see Section 2.2 for details)
Covariates: Hospital	Hospital control	Provider of services file	Hospital	Medicare Provider ID	- Categorized as for-profit, not-for-profit, government
	Registered nurse count				- Used for calculating the registered nurse per average daily census ratio. Categorized as <2, >=2 and <5, >=5
	Medical school affiliation	Impact file	Hospital	Medicare Provider ID	- Categorized as yes, no
	Transfer adjusted case mix index				- Categorized as <1.25, >=1.25 and <1.5, >=1.5
	Geographic location				- Categorized as rural (< 50,000 persons), other urban (>50,000 and <1 million), large urban (>1 million population)

Variable Category	Variable	Data Source	Level	Hospital Linkage	Specification
	Average daily census				- Used for calculating the registered nurse per average daily census ratio. Categorized as <2, >=2 and <5, >=5
	Number of beds				- Categorized as small (<200), medium (>=200 and <400), large (>=400)
	Herfindahl-Hirschman index (HHI)	Dartmouth atlas + Impact file	Hospital	Medicare Provider ID	- Categorized as unconcentrated (HHI<=1500), moderately concentrated (HHI>1500 and <=2500), highly concentrated (>2500)

Abbreviations: HAC: Hospital-Acquired Conditions; HCUP SID: Healthcare Cost and Utilization Project's State Inpatient Database; AHA ID: American Hospital Association Identification Number; DPP: Disproportionate Patient Percent; EMRAM: Electronic Medical Record Adoption Model; HIMSS: Hospital Information and Management Systems Society

2.3.2. Outcomes: Hospital-Acquired Conditions

- The following HACs were identified using the ICD-9-CM diagnosis codes specified by the Program (Table 1) that were coded as secondary diagnoses ²⁴⁵, and the POA indicators: :
 - Catheter associated bloodstream infections (BSI),
 - Catheter associated urinary tract infections (UTI),
 - Falls and trauma (FAL),
 - Poor glucose control (GLU),
 - Air embolism (AIR),
 - Blood incompatibility (BLD),
 - Surgical site infections among patients undergoing Coronary Artery Bypass Graft (CabgSSI) and spine surgery (SpineSSI), and
 - Deep vein thrombosis / pulmonary embolism among patients undergoing hip or knee replacements (HipDVT, KneeDVT).
- The POA indicators were used to determine if the secondary diagnoses of interest were present or not at admission. Stays with POA codes of N (indicating that the condition was not present at the time of admission) or U (indicating that the documentation is insufficient to determine if the condition was present at the time of admission) for HAC-related ICD-9-CM diagnosis codes were marked as stays that developed HACs ²⁴⁶. Stays with POA codes of Y (indicating that the condition was present at the time of admission) or W (clinical inability of the provider to determine the time of onset of the condition based on data and clinical judgment) for HAC-related ICD-9-CM diagnosis codes were marked as cases that did not develop HACs after admission to the hospital. The difference between the U and W codes lies in the

extent of completeness of documentation for deciding on the onset of the condition. For the U code, the documentation is incomplete and hence the hospital is subject to disincentives. For the W code, despite the completeness of the documentation, the provider is unable to decide on the onset of the condition and hence the hospital is not penalized for it.

- This approach to define the HAC outcomes slightly deviates from CMS's methodology to identify HACs that takes into consideration the reassignment of the MS-DRG. For assessing the financial disincentives, CMS considers those conditions as HACs that meet the ICD-9-CM diagnosis code and POA indicator criteria described above, and that lead to the reassignment of the MS-DRG from a lower paying to a higher paying MS-DRG. However, for this dissertation I do not use the reassignment of the MS-DRG criteria. The rationale for this approach to include all instances of HACs, regardless of whether or not the HAC led to the reassignment of the MS-DRG, is that while implementing quality improvement programs to prevent a particular HAC, hospitals are unlikely to differentiate between situations where an MS-DRG may or may not be reassigned. Furthermore, this approach used in this dissertation is similar to that used by the CMS to compute publicly reported HAC rates²⁴⁷.
- The ICD-9-CM diagnosis codes for grade III and IV pressure ulcers were introduced with the start of the HAC Program in October 2008. Because of this, the pre-Program distribution of pressure ulcers cannot be determined, and hence they are excluded from this study.
- The main analysis comprises of the following six primary outcomes:

- Any-or-none indicator for medical cohort: This is a binary indicator for each stay in the medical cohort, with 1 indicating that at least one of the six HACs (BSI, UTI, FAL, GLU, AIR, and BLD) developed during hospital stay²⁴⁸. Since the ICD-9-C codes for BSI were introduced in October 2007, calculation of BSI and this any-or-none indicator begins in October 2007 as compared to that for other HAC indicators where the pre-phase begins in January 2005.
- Any-or-none indicator for surgical cohort: This is a binary indicator for each stay in the surgical cohort, with 1 indicating that at least one of the six HACs (BSI, UTI, FAL, GLU, AIR, and BLD) developed during hospital stay²⁴⁸. The pre-phase for calculating this indicator begins in October 2007.
- CabgSSI: This is a binary indicator for occurrence of surgical site infections among patients undergoing Coronary Artery Bypass Graft.
- SpineSSI: This is a binary indicator for occurrence of surgical site infections among patients undergoing spine surgeries.
- HipDVT: This is a binary indicator for deep vein thrombosis or pulmonary embolism among patients undergoing hip replacements.
- KneeDVT: This is a binary indicator for deep vein thrombosis or pulmonary embolism among patients undergoing knee replacements.
- Computing an any-or-none indicator has reasonable validity for quantifying the underlying latent construct of safety. HACs are a relatively infrequent, yet they are unintended and undesirable outcomes of care for both patients and providers, and preventing them from occurring is a key priority. Furthermore, preventing a HAC from occurring requires a multidisciplinary systems approach. Inadequate synergy in the system is likely to contribute not only to the occurrence of the HACs of interest, but

also to other medical errors that are not under the purview of the Program. Thus, capturing the occurrence of even a single HAC using the any-or-none indicator is a rational measure to assess the overall risk of HACs, and consequently the safety environment.

- The secondary outcomes of interest in the main analysis were binary indicators for the six HACs (BSI, UTI, FAL, GLU, AIR, and BLD) in each of the cohorts.
- For the descriptive analysis, the outcomes were constructed as mean hospital-level rates by Program-phase and MUR quartile (Table 7), and by month and year of discharge (Figure 5). These rates were computed by first calculating the number of HACs per eligible stays (as defined by the cohort / sub-cohort) as a percentage to generate a hospital-level rate. This statistic was then averaged across the Program-phase and MUR quartile (Table 7), and the month and year of discharge (Figure 5) to generate the mean hospital-level HAC rate.

2.3.3. Key independent variable

The Program effect was modeled using an interaction between the Program phase (pre-/post-) and the Medicare Utilization Ratio (MUR) quartile ⁶⁵ for the hospital, which is the proportion of total inpatient days for a hospital that are attributed to Medicare. Inpatient stays with discharges from January 2005 (October 2007 for the any-or-none indicators and BSI in the medical and surgical cohorts) to September 2008 were included in the pre-Program cohort. Stays with discharges from October 2008 to December 2012 were included in the post-Program cohort.

The MUR variable was constructed as a time-constant measure, by computing its mean for the hospital during the pre-Program phase, to address the potential endogeneity of the MUR variable to the outcomes. While it is reasonable to state that hospitals with higher MUR are likely to have lower HACs following the Program, it is also possible that hospitals with lower HACs, and consequently better quality, will attract a larger proportion of Medicare patients. Fixing the value of the MUR variable over time, not only controls for the sensitivity of the variable to the introduction of the Program, but also to other quality reforms that may influence the MUR for a hospital. For example, hospitals with high HAC rate could make efforts to reduce their MUR for high-risk patients, rather than HAC rates, as the Program evolves.

2.3.4. Key covariate specifications

Operating profit margin

The Cost Reports were used to compute the operating profit margins for NY hospitals using the following formula:

Operating Profit Margin

$$= \frac{(\text{Net patient revenues} + \text{Other revenues}) - \text{Total operating expenses}}{\text{Net patient revenues} + \text{Other revenues}}$$

Disproportionate Patient Percent (DPP)

DPP was computed using the following formula:

$$DPP = \left(\frac{\text{Medicare \& Supplemental Security Income Days}}{\text{Total Medicare Days}} \right) + \left(\frac{\text{Medicaid, NonMedicare Days}}{\text{Total Patient Days}} \right)$$

A combination of the DPP (minimum of 15%) and the bed size is used to compute the Disproportionate Share Payment adjustment for a hospital. Alternate representations for the proportion of disadvantaged patients in a hospital are the percent of Medicaid patients in a hospital or the proportion of inpatient days reimbursed by Medicaid. However, using DPP has the advantage of including both elderly Medicare patients who are on Social Security Income and non-elderly patients reimbursed by Medicaid.

EMRAM stage

- The EMRAM stage variable represents the EHR staging criteria used by HIMSS. This variable was defined using a combination of the type of application (e.g.

Pharmacy Management System), and the status of this application (e.g. Live and Operational).

- The “Live and Operational” and “To be Replaced” status was used to indicate an ‘in use’ status for an application. Other status that were not considered as indicators of application usage include “Contracted/Not Yet Installed”, “Installation in Process”, “Not Automated”, “Not Reported”, “Not Yet Contracted”, and “Service Not Provided”.
- The following criteria were used to construct the EMRAM variable ^{249, 250}:
 - Stage 0: At least one of the basic EHR modules are not in use – Laboratory Information System (LIS), Radiology Information System (RIS), Pharmacy Management System (PMS)
 - Stage 1: LIS, RIS and PMS are in use
 - Stage 2: Fulfillment of Stage 1 criteria, and Clinical Data Repository (CDR) and CDSS are in use
 - Stage 3: Fulfillment of Stage 2 criteria, and Nursing Documentation, Electronic Medication Administration Record (EMAR), CDSS drug algorithms, PACS (for viewing of Radiology images outside the department) applications are in use
 - Stage 4: Stage 3, and CPOE and clinical protocols in CDSS are in use
 - Stage 5: Stage 4, and closed loop medication administration, and integration of the EMAR and radio frequency identification are in use and integrated with the CPOE and pharmacy.
- Hospitals that have achieved Stage 3 of the EMRAM model have the capabilities to fulfill CMS’ Stage 1 Meaningful Use criteria ²⁵⁰. EMRAM Stage 3 indicators and attestation to Stage 1 Meaningful Use criteria (obtained from CMS’ EHR Incentive

Program files) were used to create a categorical indicator of the hospital's EHR capabilities.

Herfindahl-Hirschman index

- The hospital- and zip code-level files were used to identify the Hospital Referral Regions (HRR) for each hospital in NY.
- The HRRs from the Dartmouth files and the average daily census from the Impact files were used to generate the Herfindahl-Hirschman index for market concentration²⁵¹
- Market share is computed using average daily census for hospitals in a given HRR. The HHI is then calculated as sum of squared market share, and multiplied by 10,000.

2.4. Statistical analysis

2.4.1. Empirical approach

Introduction

The central empirical approach is to use hierarchical logistic regression models with difference-in-difference (DID) estimation and hospital random effects.

Hierarchical models with random effects were chosen to account for the clustering of patient-level observations within hospitals. The outcomes in a hospital are a result of the prevailing structure and processes. Furthermore, the patients treated in a particular hospital are likely to have lesser variability in their characteristics and outcomes as compared to patients from another hospital. If the models were estimated without accounting for this clustering of patients within hospital, there is a potential that the standard errors might have been underestimated and this may have led to the identification of a significant association when there actually is none ²⁵².

DID estimation has been frequently used for program evaluation, and it involves the calculation of three sets of differences ²⁵³. The first is the difference in outcomes in the pre- and post-Program phase for each of the MUR groups. Doing so accounts for secular trends that may be present only during the post-Program phase, and may influence the incidence of HACs. The second difference is between the 'treatment' and 'control' groups for each of the phases, which accounts for the unobservable differences between the groups that may influence outcomes. The difference between these two sets of differences is the treatment effect, which is the post-Program change in HACs in the 'treatment' group that is over-and-above the trend extending from the pre-Program

phase, and after controlling for the trend in the 'control' group. In order to attribute the change in the HAC incidence to the Program, a key assumption underlying DID estimation is that the trends in the outcomes of the MUR groups are parallel before the start of the Program, and would have remained so had the Program not been implemented. This parallel trend assumption was checked by testing if the changes in the outcome across the groups varied over time in the pre-Program phase.

Choosing between random effects and fixed effects is a matter of debate. There is considerable literature that discusses the strengths and weaknesses of each of these methods, with specific focus on the satisfaction of assumptions, the variance-bias tradeoff, and the nature of variables that are to be estimated^{254, 255}. The choice of random effects for the multivariate models in this study was based on the compatibility of the variables to be estimated with the method chosen. Importantly, the MUR variable is fixed at its baseline for a hospital (2.3.3.). Since the MUR variable does not have within-hospital variance, using fixed effects would subsume the effect of this variable, and consequently not generate an estimate for it.

Specific Aim 1

The analyses for Specific Aim 1 are designed to identify patient- and hospital-level factors that predict the risk of a patient developing HACs during hospital stay, prior to implementation of the HAC Program.

The generic form of the models used is as follows:

Equation 1

$$\ln \frac{p(HAC_{ph} = 1)}{1 - p(HAC_{ph} = 1)} = \beta_0 + \beta_1 X_{ph} + \beta_2 Z_h + u_h$$

- HAC_{ph} : The dependent variable for each of the HAC models is a dichotomous variable representing the presence or absence of each HAC. For this variable, a value equal to 1 represents the presence of the HAC in patient p in hospital h .
- X_{ph} : This is the vector of variables for patient p in hospital h . This includes patient-level variables listed in Table 4.
- Z_h : This is the vector of variables for hospital h , and includes the hospital-level variables listed in Table 4.
- u_h : Hospital random effects

Previously, I had proposed to carry out this analysis in a two-step method - by first estimating models with only patient-level variables, and then adding hospital-level variables to understand the proportion the variance in the outcomes that is explained by hospital-level variability. However, given that the outcomes are binary in nature, comparing the variance of nested logistic models may not produce the desired results.

This is succinctly explained as follows ²⁵⁶ (text from Multilevel Analysis: Techniques and Applications, Second Edition [Page 133]):

“However, in logistic and probit regression (and many other generalized linear models), the underlying latent variable is rescaled, so the lowest-level residual variance is again $\frac{\pi^2}{3}$ or unity, respectively. Consequently, the values of the regression coefficients and higher-level variances are also rescaled, in addition to any real changes resulting from the changes in the model. These implicit scale changes make it impossible to compare regression coefficients across models, or to investigate how variance components change.”

Specific Aim 2 – Hypothesis 2.1

This Aim evaluates the overall impact of the Program across hospitals with varying MUR. The generic form of the empirical logit models is as follows:

Equation 2

$$\begin{aligned} \ln \frac{p(HAC_{pht} = 1)}{1 - p(HAC_{pht} = 1)} \\ = \beta_1 Policy_{pht} + \beta_2 \%MUR_{ht} + \beta_3 (Policy_{pht} \times \%MUR_{ht}) + \beta_4 X_{pht} \\ + \beta_5 Z_{ht} + \beta_6 Time_t + \xi_h \end{aligned}$$

- HAC_{pht} : Dichotomous variable representing the presence or absence of a particular HAC in patient p in hospital h at year t
- $Policy_{pht}$: Presence or absence of HAC Program. For discharges occurring on or after October 1, 2008, $Policy_{pht} = 1$. For discharges occurring before October 1, 2008, $Policy_{pht} = 0$
- $\%MUR_{ht}$: Percentage of inpatient Medicare discharges in a hospital. This factor serves as a proxy for the financial impact of the HAC Program. This variable is categorized into quartiles.
- $Policy_{pht} \times \%MUR_{ht}$: The key independent variable is an interaction between the variable indicating the presence or absence of the Program and the MUR categorized into quartiles. The coefficient (β_3) of this interaction term is used to test the hypothesis that the HAC Program was associated with a decline in HAC likelihood in hospitals.
- X_{pht} : These are the vector of patient covariates that have an association with the development of HACs. These covariates vary for discharges in a hospital over time.

- Z_{ht} : These are the vector of hospital covariates that vary over time.
- $Time_t$: Year fixed effects. Controlling for these factors accounts for unobserved, underlying secular trends that are associated with HACs.
- u_h : Hospital random effects

Specific Aim 2 – Hypothesis 2.2

This aim evaluates the overall impact of the Program across hospitals with varying MUR and varying financial strength. The financial strength is represented by the operating profit margin (Hypothesis 2.2.1) and the DPP (Hypothesis 2.2.2)

The generic form of these logit models is as follows:

Equation 3

$$\ln \frac{p(HAC_{pht}=1)}{1-p(HAC_{pht}=1)} = \beta_1 Policy_{pht} + \beta_2 \%MUR_{ht} + \beta_3 (Policy_{pht} \times \%MUR_{ht} \times Fin_{ht}) + \beta_4 X_{pht} + \beta_5 Z_{ht} + \beta_6 Time_t + \xi_h$$

The structure of this model is similar to that in Equation 2, with the addition of the following terms:

- Fin_{ht} : This represents the operating profit margin categorized as positive or negative (for Hypothesis 2.2.1), and the DPP categorized as above and below median (for Hypothesis 2.2.2).
- $Policy_{pht} \times \%MUR_{ht} \times Fin_{ht}$: The key independent variable is an interaction between the Program phase, the MUR quartiles, and the indicator of financial health. The coefficient (β_3) of this interaction term is used to test the hypothesis that the HAC Program was associated with a decline in HAC likelihood in hospitals for hospitals with positive operating financial margin (Hypothesis 2.2.1), and DPP below the median (Hypothesis 2.2.2).

Specific Aim 2 – Hypothesis 2.3

This aim evaluates the overall impact of the Program across hospitals with varying MUR and varying EHR capabilities, which is represented by the EMRAM Stage for a hospital.

The generic form of these models is as follows:

Equation 4

$$\begin{aligned} \ln \frac{p(HAC_{pht} = 1)}{1 - p(HAC_{pht} = 1)} \\ = \beta_1 Policy_{pht} + \beta_2 \%MCR_{ht} + \beta_3 (Policy_{pht} \times \%MCR_{ht} \times EMRAM_{ht}) \\ + \beta_4 X_{pht} + \beta_5 Z_{ht} + \beta_6 Time_t + \xi_h \end{aligned}$$

The structure of this model is similar to that in Equation 2, with the addition of the following terms:

- $EMRAM_{ht}$: This variable represents the HIMSS EMRAM score ²⁵⁷ of hospital h in year t . This variable classifies hospitals from Stage 0, which represents hospitals with minimal automation where modules such as laboratory, radiology, and pharmacy have not been implemented; to Stage 3+, which represents hospitals EHRs with at least the capabilities to meet Stage 3 Meaningful Use criteria laid down by the CMS.
- $Policy_{pht} \times \%MUR_{ht} \times EMRAM_{ht}$: The key independent variable is an interaction between the Program phase, the MUR quartiles, and the indicator of EHR capabilities. The coefficient (β_3) of this interaction term is used to test the hypothesis that the HAC Program was associated with a decline in HAC likelihood in hospitals with higher EHR capabilities.

2.4.2. Stratified and sensitivity analyses

Several stratified and sensitivity analyses were carried out to check for the consistency of the findings from the main models. The rationale for these analyses is listed below:

- Stratified analyses: The models were analyzed in sub-cohorts that were defined using the primary diagnosis (for the medical cohort), and by the key procedure (for the surgical cohort). By doing so, one can draw inferences of the Program effects for that particular sub-cohort as compared to that for the entire cohort in which the primary diagnosis / procedure is treated as a control variable.
- Short- and long-term effects: After the initial focus on reducing the likelihood of HACs, it is likely that the Program effects in hospitals may taper-off over time. This change in the Program effects was examined by redefining the post-Program phase as early post-Program phase (October 2008 to September 2010), and late post-Program phase (October 2010 to December 2012).
- Redefining the Program phases: Since the Program was implemented in 2008, it is likely that hospitals may have brought in the QI programs in advance of the Program start date, or they may have delayed the start of these programs to understand the impact of the Program on their financial resources. Hence, these set of analyses consisted of excluding discharges from 2008, and by including all the discharge from 2008 in the pre-Program phase.
- Redefining the MUR variable: The MUR variable was re-specified as a continuous variable, and in the categorical format using tertile- and quintile-based categories.
- Changes in coding practices: Changes in coding practices are known unintended consequences of P4P programs. These changes were accounted for in this study by controlling for an indicator of the percentage of stays with a HAC-related diagnosis

that were coded as POA each year (%POA). Because the %POA is HAC-specific, the models for the any-or-none indicators in each of the cohort cannot be re-estimated with the inclusion of the %POA indicator. Hence, only the results for secondary outcomes are presented for this sensitivity analysis.

2.5. Summary of key methods terms

- Dataset specifics: Administrative data from 2005 to 2012 for inpatient stays in NY hospitals reimbursed by IPPS.
- Study observations: Inpatient stays for elderly (≥ 65 years and ≤ 99 years) Medicare beneficiaries whose stays are reimbursed by fee-for-service Medicare.
- Medical cohort: Comprises of inpatient stays with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke
- Surgical cohort: Comprises of inpatient stays undergoing CABG, spine surgeries, hip or knee replacement.
- Primary outcomes: These include any-or none indicator (medical cohort), any-or-none indicator (surgical cohort), CabgSSI (CABG sub-cohort of surgical cohort), SpineSSI (Spine surgery sub-cohort of surgical cohort), HipDVT (Hip replacement sub-cohort of surgical cohort), and KneeDVT (Knee replacement sub-cohort of surgical cohort).
- Secondary outcomes: These include BSI, UTI, FAL, GLU, AIR, and BLD in each of the cohorts (medical and surgical).
- Main analysis: Hierarchical logistic regression models with DID estimation and hospital random effects for the primary (results presented in the main text of the monograph) and secondary (results presented in the appendix) outcomes.

Chapter 3 - RESULTS

This section presents the descriptive results and the multivariate results for the main, stratified, and sensitivity analyses. In the main and sensitivity analyses, I have presented results for the medical and surgical any-or-none indicator, HipDVT, KneeDVT, and SpineSSI. The results from the multivariate secondary analyses for BSI, UTI, FAL, and GLU for each cohort are presented in the Appendix.

3.1. Descriptive statistics

3.1.1. Hospital and patient characteristics

Descriptive statistics for hospitals and inpatient stays by the MUR quartile in each of the cohorts from 2005 to 2012 are presented in Table 5 and Table 6 respectively.

Characteristics of hospitals:

Of the 159 hospitals in the medical cohort (Table 5), 87% were non-government, 47% were small (<200 beds), 70% were located in large urban areas, 60% were affiliated to medical schools, 60% were located in competitive markets ($HHI \leq 1500$), and 48% had EHRs that were EMRAM Stage 3 or higher. In this cohort, the mean DPP was 36.11 (Standard deviation [SD]: 25.28), the mean operating profit margin was -0.01 (SD: 0.08), the mean transfer-adjusted case mix index was 1.40 (SD: 0.25), and the mean registered nurse per average daily census ratio was 2.37 (SD: 1.73). The surgical cohort comprised of 153 hospitals with similar distribution of characteristics.

Characteristics of inpatient stays:

Among the 867,584 inpatient stays of the medical cohort (Table 6), 37% had a primary diagnosis of CHF, the mean age was 80.02 years (SD: 8.19), and the mean number of comorbidities was 2.91 (SD: 1.59). In this cohort of inpatient stays, 54% were female, 68% were White non-Hispanic, 95% were emergency or urgent admissions, 90% were admitted from home or the doctor's office, 23% were admitted on the weekend, Medicare was the primary payer for 96% of the stays, and 14% underwent a major surgery.

Among the 264,713 inpatient stays of the surgical cohort (Table 6), 38% of the stays underwent knee replacements, the mean age was 75.31 years (SD: 6.96), and the mean number of comorbidities was 2.19 (SD: 1.42). In this cohort of inpatient stays, 59% were female, 78% were White non-Hispanic, 27% were emergency or urgent admissions, 92% were admitted from home or the doctor's office, 6% were admitted on the weekend, and Medicare was the primary payer for 92% of the stays.

Table 5: Descriptive statistics for hospitals in the medical and surgical cohorts

Cohort ¹	Medical					Surgical				
Variable / MUR ² Quartile	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total
n	37	41	41	40	159	32	40	41	40	153
MUR ⁶	25.37 (7.47)	38.21 (7.26)	49.53 (4.80)	58.79 (6.31)	43.32 (13.95)	25.39 (7.65)	38.28 (7.31)	49.53 (4.87)	58.64 (6.68)	43.92 (13.77)
Hospital Control (%) ³										
Not-for-profit	59.46	92.68	95.12	100.00	87.42	59.38	92.50	95.12	100.00	88.24
Government	40.54	4.88	0.00	0.00	10.69	40.62	5.00	0.00	0.00	9.80
For-profit	0.00	2.44	4.88	0.00	1.89	0.00	2.50	4.88	0.00	1.96
Number of beds (%)										
Small (<200)	29.73	26.83	56.10	75.00	47.17	18.75	25.00	56.10	75.00	45.10
Medium (>=200 & <400)	43.24	31.71	39.02	22.50	33.96	50.00	32.50	39.02	22.50	35.29
Large (>=400)	27.03	41.46	4.88	2.50	18.87	31.25	42.50	4.88	2.50	19.61
Geographic location (%)										
Rural	2.70	4.88	7.32	17.50	8.18	0.00	2.50	7.32	17.50	7.19
Large urban	91.89	87.80	68.29	32.50	69.81	96.88	90.00	68.29	32.50	70.59
Other urban	5.41	7.32	24.39	50.00	22.01	3.12	7.50	24.39	50.00	22.22
Medical school affiliation (%)	94.59	73.17	41.46	35.00	60.38	100.00	75.00	41.46	35.00	60.78
Herfindahl-Hirschman Index (%)										
Unconcentrated	72.97	63.41	63.41	42.50	60.38	75.00	65.00	63.41	42.50	60.78
Moderately concentrated	10.81	21.95	21.95	30.00	21.38	9.38	22.50	21.95	30.00	21.57
Highly concentrated	16.22	14.63	14.63	27.50	18.24	15.62	12.50	14.63	27.50	17.65

Cohort ¹	Medical					Surgical				
Variable / MUR ² Quartile	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total
EHR ⁴ capabilities (%)										
EMRAM ⁵ Stage 0	0.00	4.88	0.00	0.00	1.26	0.00	2.50	0.00	0.00	0.65
EMRAM Stage 1	5.41	0.00	2.44	15.00	5.66	3.12	0.00	2.44	12.50	4.58
EMRAM Stage 2	13.51	29.27	34.15	37.50	28.93	12.50	30.00	34.15	37.50	29.41
EMRAM Stage 3+	78.38	63.41	56.10	47.50	61.01	81.25	65.00	56.10	47.50	61.44
Disproportionate patient percent ⁶	67.22 (26.78)	38.94 (15.81)	23.19 (13.10)	17.69 (6.65)	36.11 (25.28)	67.76 (25.87)	39.02 (15.82)	23.10 (13.08)	17.80 (6.51)	35.22 (24.55)
Operating profit margin ⁶	-0.07 (0.10)	0.00 (0.06)	0.00 (0.09)	0.01 (0.06)	-0.01 (0.08)	-0.07 (0.10)	0.00 (0.06)	0.00 (0.10)	0.01 (0.06)	-0.01 (0.09)
Transfer-adjusted case mix index ⁶	1.41 (0.23)	1.49 (0.29)	1.38 (0.20)	1.32 (0.24)	1.40 (0.25)	1.46 (0.20)	1.51 (0.27)	1.38 (0.21)	1.32 (0.23)	1.41 (0.24)
Registered nurse per average daily census ratio ⁶	2.31 (1.09)	1.98 (1.21)	2.26 (1.32)	2.94 (2.68)	2.37 (1.73)	2.14 (0.64)	1.87 (0.77)	2.26 (1.34)	2.93 (2.71)	2.31 (1.66)

¹ The medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. The surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

² Medicare Utilization Ratio – This is the percent of inpatient days in a hospital that are attributed to Medicare.

³ % refers to percentage of total.

⁴ Electronic Health Record

⁵ Electronic Medical Record Adoption Model

⁶ Numbers indicate mean with standard deviation in parentheses

Table 6: Descriptive statistics for inpatient stays in the medical and surgical cohorts

Cohort¹	Medical					Surgical				
Variable / MUR² Quartile	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total
n	187,321	322,096	192,222	165,945	867,584	39,150	95,870	79,221	50,472	264,713
Primary diagnosis (%) ³										
Acute myocardial infarction	17.71	18.05	15.19	16.35	17.02	Not applicable				
Congestive heart failure	38.69	36.65	37.78	36.61	37.33					
Pneumonia	21.38	23.40	28.62	29.34	25.25					
Stroke	22.23	21.91	18.41	17.70	20.40					
Surgery of interest (%)										
CABG	Not applicable					22.99	30.05	7.93	21.26	20.71
Spine						10.35	9.45	7.91	4.60	8.20
Hip replacement						30.51	29.75	38.96	34.56	33.54
Knee replacement						36.16	30.75	45.20	39.57	37.56
Age in years: Mean (SD) ⁴	78.67 (8.24)	79.92 (8.18)	80.94 (8.10)	80.66 (8.02)	80.02 (8.19)	74.60 (6.84)	75.54 (7.00)	75.08 (6.89)	75.78 (7.03)	75.31 (6.96)
Female (%)	55.67	53.44	55.44	53.71	54.42	60.52	56.99	62.33	58.57	59.41
Race & Ethnicity (%)										
White, non-Hispanic	43.06	69.04	73.63	88.39	68.15	67.23	75.35	80.72	86.82	77.94
Black, non-Hispanic	26.59	11.52	7.91	3.14	12.37	12.07	6.80	3.82	1.89	5.75
Hispanic	14.90	8.38	5.29	3.64	8.20	8.46	5.37	4.26	3.89	5.21

Cohort¹	Medical					Surgical				
Variable / MUR² Quartile	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total	1 (<35.25)	2 (>35.25& <=46.64)	3 (>46.64& <=56.86)	4 (>56.86)	Total
Emergent/Urgent admission (%)	96.35	93.39	95.37	93.75	94.54	29.67	32.64	16.95	28.70	26.75
Admitted from home (%)	91.50	87.62	91.67	91.05	90.01	87.94	89.20	97.63	93.96	92.45
Admission on weekend (%)	23.70	22.67	24.10	23.44	23.36	5.66	6.11	4.91	5.50	5.57
Primary payer: Medicare (%)	95.67	95.56	95.74	96.85	95.87	88.95	91.66	91.41	93.89	91.61
Major surgical procedure (%)	13.12	18.98	9.14	12.76	14.34	Not applicable				
Median income quartile for zip code (%)										
1	37.49	19.98	17.49	15.38	22.33	24.60	14.40	12.00	14.64	15.23
2	20.95	20.78	25.16	25.82	22.75	24.35	20.20	22.06	27.23	22.71
3	17.85	29.24	24.03	25.25	24.86	24.15	29.34	24.73	26.08	26.57
4	11.87	29.16	31.42	31.88	26.45	23.07	34.89	39.58	30.23	33.66
Comorbidities ^b : Mean (SD)	2.88 (1.54)	2.80 (1.56)	3.01 (1.63)	3.06 (1.66)	2.91 (1.59)	2.23 (1.41)	2.20 (1.40)	2.12 (1.42)	2.27 (1.46)	2.19 (1.42)
Hypertension	75.81	71.32	69.45	67.28	71.11	74.56	72.11	69.40	73.06	71.84
Chronic pulmonary disease	28.29	28.16	34.92	37.52	31.48	17.38	15.61	14.53	18.54	16.11
Diabetes mellitus, uncomplicated	34.98	29.66	28.53	27.50	30.14	22.99	22.21	17.13	21.72	20.71
Fluid & electrolyte disorder	24.21	24.11	26.26	28.28	25.40	14.91	16.79	11.84	17.05	15.08
Hypothyroidism	11.53	13.64	15.73	15.86	14.07	13.26	13.78	16.14	14.96	14.63

¹ The medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. The surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

² Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

³ % refers to percentage of total.

⁴ Numbers indicate mean with standard deviation in parentheses

⁵ Statistics for the five most frequently occurring comorbidities are listed in this table. The multivariate models include 29 indicators for each of the comorbidities of the modified Elixhauser algorithm.

3.1.2. Hospital-Acquired Conditions

Figure 5 presents Lowess curves for the overall unadjusted hospital-level mean monthly rates (Y-axis) for six primary outcomes over time (X-axis). The medical any-or-none indicator (subgraph A), and the surgical any-or-none indicator (subgraph B) show a distinct decline in their respective rates over time. The curves for CabgSSI (subgraph C), SpineSSI (subgraph D), and HipDVT (subgraph E) appear to be relatively stable over the two study phases. The curve for KneeDVT (subgraph F) appears to have a slight incline immediately after the start of the Program, and this is then followed by a decline.

Figure 5: Lowess graphs for unadjusted mean rates for primary Hospital-Acquired Condition outcomes

Table 7 provides the unadjusted hospital-level mean HAC rates (HACs per 100 eligible stays in a hospital) for the primary outcomes by the Program phase and MUR quartile. Table A 1 and Table A 2 of the Appendix provides the unadjusted hospital-level mean HAC rates for each individual HAC in the medical and surgical cohort respectively. In the medical cohort, the mean hospital-level rate for the any-or-none indicator declined 0.04 percentage points (from 0.23% [SD: 0.31] in the pre-phase to 0.19% [SD: 0.18] in the post-phase) over the study period. The number of hospitals reporting at least one HAC increased from 58% in the pre-phase to 82% in the post-phase. In both phases, hospitals in quartile 2 had the highest mean any-or-none HAC rate (0.34% [SD: 0.37] in the pre-phase, and 0.22% [SD: 0.25] in the post-phase), and the highest number of hospitals reporting at least one HAC (82% in the pre-phase, and 89% in the post-phase). In the surgical cohort, the mean hospital-level rate for the any-or-none indicator declined 0.49 percentage points (from 0.85% [SD: 3.01] in the pre-phase to 0.36% [SD: 0.52] in the post-phase) over the study period. The number of hospitals reporting at least one HAC increased from 46% in the pre-phase to 70% in the post-phase. In the pre-phase, hospitals in quartile 2 had the highest mean any-or-none HAC rate (1.54% [SD: 5.30]), and in the post-phase hospitals in quartile 3 had the highest mean any-or-none HAC rate (0.46% [SD: 0.63]). In both phases, quartile 2 had the highest number of hospitals reporting at least one HAC (67% in the pre-phase, and 86% in the post-phase).

Among the other primary outcomes, the unadjusted mean SpineSSI rate declined by 0.48 percent points (1.59% in the pre-phase, and 1.11% in the post-phase) in the post-Program phase. However, the mean rates for CabgSSI increased by 0.02 percent points (from 0.05% in the pre-phase to 0.07% in the post-phase), for HipDVT increased by 0.04

percent points (from 0.82% in the pre-phase to 0.86% in the post-phase), and for KneeDVT increased by 0.31 percent points (from 0.64% in the pre-phase to 0.95% in the post-phase) in the post-Program phase.

.

Table 7: Unadjusted hospital-level Hospital-Acquired Condition rates in the cohort¹

Program Phase	Pre ²					Post ³				
MUR ⁴ Quartile	1	2	3	4	Total	1	2	3	4	Total
Medical cohort: Any-or-none⁵										
Hospitals (inpatient stays)	35 (22,934)	39 (39,297)	41 (24,232)	38 (20,532)	153 (106,995)	34 (92,224)	38 (161,676)	39 (92,641)	37 (78,523)	148 (425,064)
Mean rate (SD) ⁶	0.14 (0.18)	0.34 (0.37)	0.20 (0.31)	0.23 (0.30)	0.23 (0.31)	0.16 (0.11)	0.22 (0.25)	0.17 (0.14)	0.20 (0.18)	0.19 (0.18)
Hospitals with at least 1 HAC (%) ⁷	45.71	82.05	48.78	52.63	57.52	85.29	89.47	79.49	75.68	82.43
Surgical cohort: Any-or-none⁵										
Hospitals (inpatient stays)	30 (4,858)	39 (11,379)	41 (9,691)	36 (6,371)	146 (32,299)	31 (19,154)	37 (51,812)	39 (44,840)	34 (25,521)	141 (141,327)
Mean rate (SD)	1.26 (2.35)	1.54 (5.30)	0.26 (0.48)	0.45 (1.02)	0.85 (3.01)	0.27 (0.41)	0.37 (0.40)	0.46 (0.63)	0.32 (0.58)	0.36 (0.52)
Hospitals with at least 1 HAC (%)	46.67	66.67	34.15	36.11	45.89	48.39	86.49	82.05	58.82	70.21
Coronary Artery Bypass Graft: Surgical Site Infections (CabgSSI)										
Hospitals (inpatient stays)	9 (5,294)	20 (14,939)	6 (3,453)	4 (5,659)	39 (29,345)	9 (3,705)	18 (13,872)	5 (2,828)	5 (5,071)	37 (25,476)
Mean rate (SD)	0.03 (0.07)	0.06 (0.14)	0.07 (0.17)	0.00 (0.00)	0.05 (0.12)	0.00 (0.00)	0.08 (0.11)	0.27 (0.61)	0.00 (0.00)	0.07 (0.23)
Hospitals with at least 1 CabgSSI (%)	22.22	20.00	16.67	0.00	17.95	0.00	38.89	20.00	0.00	21.62
Spine Surgeries: Surgical Site Infections (SpineSSI)										
Hospitals (inpatient stays)	37 (1,900)	33 (3,164)	27 (2,267)	25 (1,011)	112 (8,342)	29 (2,151)	31 (5,895)	23 (3,996)	20 (1,312)	103 (13,354)

Program Phase	Pre ²					Post ³				
MUR ⁴ Quartile	1	2	3	4	Total	1	2	3	4	Total
Mean rate (SD)	0.32 (0.58)	1.56 (3.40)	0.41 (0.89)	4.28 (19.97)	1.59 (9.60)	2.69 (9.25)	0.67 (1.08)	0.38 (0.84)	0.32 (0.89)	1.11 (5.01)
Hospitals with at least 1 SpineSSI (%)	29.63	45.45	22.22	12.00	28.57	41.38	51.61	26.09	15.00	35.92
Hip Replacement: Deep vein thrombosis / pulmonary embolism (HipDVT)										
Hospitals (inpatient stays)	32 (5,899)	40 (12,979)	41 (13,460)	39 (8,600)	152 (40,938)	31 (6,045)	37 (15,545)	39 (17,408)	34 (8,845)	141 (47,843)
Mean rate (SD)	1.36 (2.88)	0.91 (0.95)	0.61 (0.76)	0.48 (0.77)	0.82 (1.53)	1.47 (1.78)	1.03 (1.09)	0.48 (0.66)	0.53 (1.01)	0.86 (1.22)
Hospitals with at least 1 HipDVT (%)	50.00	72.50	58.54	46.15	57.24	64.52	75.68	56.41	50.00	61.70
Knee Replacement: Deep vein thrombosis / pulmonary embolism (KneeDVT)										
Hospitals (inpatient stays)	29 (6,903)	40 (12,976)	41 (15,201)	40 (9,681)	150 (44,761)	30 (7,253)	36 (16,500)	39 (20,608)	34(10,293)	139 (54,654)
Mean rate (SD)	0.51 (1.00)	0.95 (1.39)	0.64 (1.15)	0.42 (0.83)	0.64 (1.13)	1.07 (1.69)	1.41 (2.14)	0.58 (0.97)	0.78 (2.29)	0.95 (1.84)
Hospitals with at least 1 KneeDVT (%)	41.38	65.00	48.78	45.00	50.67	50.00	75.00	53.85	55.88	58.99

¹ Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

² Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries.

³ Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁴ *Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.*

⁵ *Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility)*

⁶ *Numbers indicate mean hospital-level HAC rate with standard deviation in parentheses. The HAC rate is computed as a percentage.*

⁷ *Numbers indicate percentage of hospitals with at least one HAC.*

3.2. Specific Aim 1

The analyses for Specific Aim 1 are designed to identify patient- and hospital-level factors that predict the risk of a patient developing HACs during hospital stay, prior to implementation of the HAC Program. Additionally, the analyses aid in understanding the extent of variation in the risk of HACs that can be attributed to each level of the hierarchical data. Table 8 contains the odds ratios for the any-or-none indicators in each cohort, and for SpineSSI, HipDVT, and KneeDVT outcomes. The model for CabgSSI outcome did not converge, and hence this outcome is excluded from this table and from the subsequent analyses.

Medical Cohort – Any-or-none indicator:

After controlling for risk factors and hospital random effects, the relative odds of developing at least one HAC (Table 8 – Column 2) in the medical cohort in the pre-Program phase were significantly higher among (i) MUR 2 hospitals (Conditional Odds Ratio [COR]: 1.73; 95% CI [Confidence Interval]: 1.08 to 2.80) as compared to MUR 1 hospitals; (ii) stays with major surgeries (COR: 3.01; 95% CI: 2.11 to 4.29) as compared to those with minor surgeries; and (iii) those with certain comorbidities. The relative odds were significantly lower among (i) those with elective admissions (COR: 0.20; 95% CI: 0.08 to 0.46) as compared to those with emergency or urgent admissions; (ii) Black non-Hispanics (COR: 0.50; 95% CI: 0.32 to 0.78) as compared to White non-Hispanics; (iii) those without surgeries as compared to those with minor surgeries (COR: 0.32, 95% CI: 0.22 to 0.47); and (iv) among those with certain comorbidities.

Surgical Cohort:

After controlling for risk factors and hospital random effects, the relative odds of developing at least one HAC (Table 8 – Column 3) in the surgical cohort in the pre-Program phase were significantly higher for those with certain comorbidities. The relative odds were significantly lower among (i) those with elective admissions (COR: 0.41; 95% CI: 0.27 to 0.63) as compared to those with emergency or urgent admissions; and (ii) among those treated in large hospitals (COR: 0.36; 95% CI: 0.16 to 0.82) as compared to those treated in small hospitals; and (iii) among those with certain comorbidities.

For SpineSSI Table 8 – Column 4), discharges in 2008 (COR: 2.49; 95% CI: 1.06 to 5.86), private primary payer (COR: 1.74; 95% CI: 1.06 to 2.87), and MUR quartile 2 (COR: 2.07; 95% CI: 1.02 to 4.21) were predictors of increased risk. Patient age ≥ 70 and < 80 years (COR: 0.60; 95% CI: 0.37 to 0.98), elective admissions (COR: 0.34; 95% CI: 0.18 to 0.64), Stage 2 EMRAM hospitals (COR: 0.29; 95% CI: 0.09 to 0.95), and certain comorbidities were predictive of decreased risk in the pre-Program phase.

For HipDVT, age ≥ 70 and < 80 years (COR: 1.36; 95% CI: 1.03 to 1.79), female gender (COR: 1.30; 95% CI: 1.03 to 1.64), certain comorbidities, and MUR quartile 2 hospitals (COR: 1.57; 95% CI: 1.02 to 2.41) were predictive of increased risk in the pre-Program phase. Stays transferred from other hospitals (COR: 0.34; 95% CI: 0.14 to 0.80), certain comorbidities, moderately concentrated hospitals (COR: 0.58; 95% CI: 0.37 to 0.91), and hospitals with RN/ADC ratio of ≥ 5 (COR: 0.52; 95% CI: 0.28 to 0.95) were predictive of decreased risk in the pre-Program phase.

For KneeDVT, certain comorbidities were predictive of decreased risk in the pre-Program phase, whereas discharge years of 2006 (COR: 0.69; 95% CI: 0.49 to 0.97) and 2007 (COR: 0.51; 95% CI: 0.31 to 0.83), and certain comorbidities were predictive of decreased risk.

Importantly, the hospital-level variance ranged from 0.15 for the surgical any-or-none indicator to 0.86 for KneeDVT, while the intraclass correlation coefficient ranged from 1% for the surgical any-or-none indicator to 18% for KneeDVT. Based on the chi-square statistic for the likelihood ratio test for clustering, hospital-level variability is significant for the medical any-or-none indicators, HipDVT, and KneeDVT in the pre-Program phase.

Table 8: Odds ratios of risk factors for the any-or-none indicator in the pre-Program phase¹

Cohort²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none³</i>	<i>Any-or-none</i>	<i>SpineSSI⁴</i>	<i>HipDVT⁵</i>	<i>KneeDVT⁶</i>
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Medicare Utilization Ratio quartile					
1: <=35.25%	Ref	Ref	Ref	Ref	Ref
2: >35.25% & <=46.64%	1.73* [1.08,2.80]	1.39 [0.68,2.84]	2.07* [1.02,4.21]	1.57* [1.02,2.41]	1.56 [0.56,4.34]
3: >46.64% & <=56.86%	1.03 [0.50,2.13]	0.42 [0.17,1.05]	2.38 [0.87,6.53]	1.61 [0.96,2.72]	2.17 [0.04,111.52]
4: >56.86% & <=80.33%	1.22 [0.61,2.47]	0.51 [0.19,1.37]	2.29 [0.58,9.08]	1.57 [0.86,2.89]	1.83 [0.36,9.31]
Discharge year					
2005			Ref	Ref	Ref
2006			1.15 [0.51,2.62]	1.01 [0.66,1.57]	0.69* [0.49,0.97]
2007	Ref	Ref	1.82 [0.75,4.41]	0.79 [0.47,1.33]	0.51** [0.31,0.83]
2008	1.12 [0.82,1.52]	1.22 [0.75,1.98]	2.49* [1.06,5.86]	0.95 [0.53,1.72]	0.61 [0.30,1.20]
Age in years					
<70	Ref	Ref	Ref	Ref	Ref
>=70 & <80	0.92 [0.62,1.37]	0.83 [0.55,1.25]	0.60* [0.37,0.98]	1.36* [1.03,1.79]	0.99 [0.83,1.20]
>=80 & <90	1.07 [0.75,1.53]	1.00 [0.61,1.65]	0.69 [0.35,1.35]	1.26 [0.90,1.76]	0.82 [0.56,1.20]
>=90	1.28 [0.82,1.99]	0.70 [0.34,1.44]	3.85 [0.42,35.29]	1.18 [0.76,1.83]	1.00 [1.00,1.00]
Admission type					
Emergent/Urgent/Trauma	Ref	Ref	Ref	Ref	Ref
Elective	0.20*** [0.08,0.46]	0.41*** [0.27,0.63]	0.34*** [0.18,0.64]	1.02 [0.74,1.41]	0.69 [0.38,1.28]
Gender					
Male	Ref	Ref	Ref	Ref	Ref
Female	1.32 [0.99,1.75]	1.08 [0.76,1.55]	1.34 [0.75,2.38]	1.30* [1.03,1.64]	1.06 [0.70,1.62]
Race and ethnicity					

Cohort²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none³</i>	<i>Any-or-none</i>	<i>SpineSSI⁴</i>	<i>HipDVT⁵</i>	<i>KneeDVT⁶</i>
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
White, non-Hispanic	Ref	Ref	Ref	Ref	Ref
Black, non-Hispanic	0.50** [0.32,0.78]	0.90 [0.42,1.93]	0.74 [0.26,2.11]	0.72 [0.36,1.44]	1.31 [0.97,1.76]
Hispanic	0.77 [0.52,1.13]	1.41 [0.71,2.78]	0.39 [0.09,1.70]	1.24 [0.72,2.13]	0.96 [0.63,1.45]
Other, non-Hispanic	0.92 [0.65,1.31]	1.37 [0.65,2.90]	0.77 [0.40,1.46]	0.88 [0.46,1.71]	1.08 [0.57,2.04]
Race miss/Eth unknown	1.06 [0.54,2.07]	1.29 [0.63,2.66]	0.79 [0.35,1.81]	1.49* [1.03,2.15]	1.27 [0.84,1.92]
Primary payer					
Medicare	Ref	Ref	Ref	Ref	Ref
Private	0.98 [0.54,1.78]	0.83 [0.39,1.76]	1.74* [1.06,2.87]	1.34 [0.83,2.17]	1.06 [0.67,1.68]
Other/unins/self/Medicaid	1.91 [0.70,5.21]	4.30*** [2.05,9.04]	4.50** [1.51,13.45]	0.86 [0.08,9.15]	1.98 [0.98,3.98]
Transfer-in					
No	Ref	Ref	Ref	Ref	Ref
Yes	0.91 [0.58,1.42]	0.83 [0.47,1.47]	0.94 [0.39,2.28]	0.34* [0.14,0.80]	0.64 [0.04,9.76]
EMRAM stage					
Stage 0	Ref	Ref	Ref	Ref	Ref
Stage 1	0.64 [0.15,2.76]	1.23 [0.22,6.77]	0.40 [0.11,1.48]	0.96 [0.56,1.62]	2.06 [0.51,8.25]
Stage 2	0.52 [0.13,2.16]	0.53 [0.10,2.83]	0.29* [0.09,0.95]	1.01 [0.64,1.60]	1.23 [0.50,3.04]
HHI (Concentration)					
Unconcentrated	Ref	Ref	Ref	Ref	Ref
Moderately concentrated	0.78 [0.52,1.18]	0.72 [0.42,1.24]	0.52 [0.23,1.20]	0.58* [0.37,0.91]	1.66 [0.87,3.18]
Highly concentrated	1.12 [0.55,2.29]	1.20 [0.59,2.44]	0.33 [0.07,1.53]	0.79 [0.45,1.39]	0.59 [0.15,2.32]
RN/ADC Ratio					
<2	Ref	Ref	Ref	Ref	Ref
>=2 & <5	0.90 [0.61,1.34]	1.05 [0.62,1.76]	1.80 [0.76,4.24]	0.75 [0.52,1.09]	0.75 [0.44,1.27]
>=5	1.20 [0.33,4.36]	2.78 [0.89,8.70]	1.61 [0.41,6.36]	0.52* [0.28,0.95]	0.51 [0.16,1.65]

Cohort²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none³</i>	<i>Any-or-none</i>	<i>SpineSSI⁴</i>	<i>HipDVT⁵</i>	<i>KneeDVT⁶</i>
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Surgical procedure					
Minor	Ref				
Major	3.01*** [2.11,4.29]				
None	0.32*** [0.22,0.47]				
Elixhauser comorbidities					
Iron deficiency anemia	0.72 [0.50,1.03]	1.25 [0.83,1.88]	1.18 [0.49,2.83]	1.24 [0.92,1.67]	1.58*** [1.22,2.03]
Chronic blood loss anemia	1.37 [0.50,3.74]	0.54 [0.07,4.18]	1.00 [1.00,1.00]	1.64 [0.68,3.95]	2.57** [1.34,4.92]
Congestive heart failure	1.75** [1.18,2.61]	1.95** [1.21,3.12]	2.06 [0.90,4.70]	1.41 [0.97,2.04]	1.32 [0.85,2.05]
Chronic pulmonary disease	0.89 [0.69,1.14]	0.95 [0.65,1.39]	1.25 [0.75,2.11]	1.32* [1.03,1.71]	1.03 [0.65,1.63]
Coagulopathy	1.67* [1.02,2.74]	1.05 [0.52,2.15]	0.71 [0.20,2.47]	1.67* [1.00,2.77]	2.30* [1.07,4.94]
Diabetes mellitus (uncomplicated)	0.92 [0.67,1.26]	0.97 [0.54,1.71]	0.92 [0.43,1.95]	0.67* [0.45,0.99]	1.06 [0.81,1.39]
Drug abuse	5.36** [1.55,18.46]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.14 [0.11,11.45]
Hypertension	0.58*** [0.46,0.73]	0.45*** [0.32,0.62]	0.43*** [0.29,0.65]	0.70** [0.57,0.88]	0.77** [0.64,0.92]
Hypothyroidism	0.62* [0.39,1.00]	0.60 [0.32,1.10]	0.73 [0.29,1.87]	0.78 [0.56,1.07]	0.82 [0.58,1.16]
Lymphoma	1.63 [0.72,3.67]	1.00 [1.00,1.00]	3.13 [0.68,14.42]	2.78** [1.38,5.58]	2.20 [0.21,22.92]
Fluid and electrolyte disorder	1.45** [1.12,1.87]	1.47 [0.97,2.23]	1.62 [0.99,2.63]	1.95*** [1.41,2.68]	1.51* [1.01,2.25]
Metastatic cancer	0.82 [0.33,2.05]	1.12 [0.28,4.46]	0.84 [0.20,3.51]	3.55** [1.58,7.98]	2.65 [0.46,15.10]
Obesity	0.81 [0.43,1.53]	0.67 [0.26,1.74]	1.11 [0.42,2.94]	1.60* [1.06,2.42]	1.13 [0.84,1.51]
Paralysis	2.21*** [1.40,3.50]	1.93 [0.85,4.40]	0.87 [0.34,2.22]	1.02 [0.34,3.04]	0.91 [0.23,3.53]
Pulmonary circulation disorder	1.19 [0.53,2.65]	1.54 [0.68,3.50]	0.75 [0.08,6.72]	23.18*** [15.99,33.60]	24.59 [0.40,1494.55]
Renal failure	1.38* [1.00,1.89]	1.04 [0.64,1.70]	1.96 [0.74,5.16]	1.87** [1.24,2.83]	1.33 [0.77,2.31]
Solid tumor without metastasis	1.37 [0.75,2.52]	0.75 [0.20,2.82]	2.83 [0.75,10.66]	1.97* [1.17,3.29]	0.85 [0.14,5.25]

Cohort²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none³</i>	<i>Any-or-none</i>	<i>SpineSSI⁴</i>	<i>HipDVT⁵</i>	<i>KneeDVT⁶</i>
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Valvular disease	0.95 [0.57,1.59]	1.01 [0.59,1.70]	1.19 [0.64,2.22]	0.41*** [0.27,0.64]	0.54 [0.17,1.79]
Weight loss	1.94** [1.18,3.17]	3.10** [1.51,6.35]	1.80 [0.52,6.25]	1.32 [0.58,3.00]	1.43 [0.33,6.23]
Number of beds					
Small (<200)	Ref	Ref	Ref	Ref	Ref
Medium (>=200 & <400)	0.94 [0.51,1.73]	0.63 [0.33,1.18]	1.53 [0.57,4.10]	1.01 [0.65,1.58]	1.29 [0.05,35.29]
Large (>=400)	0.86 [0.42,1.76]	0.36* [0.16,0.82]	1.17 [0.55,2.49]	2.14** [1.21,3.80]	3.17 [0.04,226.65]
Model parameters					
Inpatient observations	106,400	32,071	8,223	40,725	44,363
Hospitals	151	145	109	151	149
Hospital-level random effect	0.56	0.15	0.45	0.38	0.86
Intraclass correlation coefficient	0.09	0.01	0.06	0.04	0.18
Likelihood ratio test					
Chi-square statistic	17.01	0.06	2.02	10.64	111.61
p-value	<0.001	0.40	0.08	<0.001	<0.001

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group; EMRAM: Electronic Medical Record Adoption Model; HHI: Herfindahl-Hirschman Index; RN/ADC: Registered nurse per average daily census ratio

¹ Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

² Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and from January 2005 to September 2008 for SpineSSI, HipDVT and KneeDVT.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ *SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.*

⁵ *HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.*

3.3. Specific Aim 2.1

This Aim evaluates the overall impact of the Program across hospitals with varying MUR.

3.3.1. Hypothesis 2.1

Controlling for relevant variables, the HAC Program is associated with a decline in the HAC likelihood among Medicare beneficiaries admitted to NY hospitals, and that this decline is greater among hospitals at a higher risk of financial losses from the Program.

Table 9 presents the results from the hierarchical logistic regression models for this Aim. Figure 6 graphically represents the multivariate odds ratios tabulated in Table 9. Appendix Table A 3 presents the results of this analysis on the secondary outcomes in each cohort. The interpretation of the main and interaction effects is as follows²⁵⁸:

Program main effects:

After controlling for patient factors, hospital factors, and hospital random effects; for MUR Quartile 1 hospitals, the Program was not associated with significant changes in the relative odds of the five primary outcomes investigated.

MUR main effects:

From the multivariate analyses presented in Table 9, in the pre-Program phase, stays in MUR Quartile 2 hospitals had significantly higher relative odds of medical any-or-none indicator (COR: 1.55; 95% CI: 1.03 to 2.34), of SpineSSI (COR: 2.38; 95% CI: 1.13 to 5.00), and of HipDVT (COR: 1.98; 95% CI: 1.28 to 3.06) as compared to MUR Quartile 1

hospitals. Furthermore, for HipDVT, stays in MUR Quartile 3 (COR: 1.84; 95% CI: 1.13 to 2.99) and MUR Quartile 4 (COR: 1.82; 95% CI: 1.05 to 3.15) hospitals had a significantly higher relative odds of HipDVT as compared to MUR Quartile 1 hospitals.

Program and MUR interaction effects:

For the medical cohort, after controlling for patient factors, hospital factors, and hospital random effects, the Program was associated with a significant decline in the likelihood of the any-or-none indicator among MUR quartile 2 hospitals as compared to MUR quartile 1 hospitals (COR: 0.57; 95% CI: 0.38 to 0.87). Furthermore, the Program was associated with a significant decline in the likelihood of SpineSSI (for Quartile 2 – COR: 0.34, 95% CI: 0.14 to 0.82; for Quartile 3 – COR: 0.41, 95% CI: 0.18 to 0.94), HipDVT (for Quartile 2 – COR: 0.56, 95% CI: 0.34 to 0.91; for Quartile 3 – COR: 0.46, 95% CI: 0.27 to 0.79; for Quartile 4 – COR: 0.51, 95% CI: 0.26 to 1.00), and KneeDVT (for Quartile 3 – COR: 0.36, 95% CI: 0.18 to 0.74). Overall, the Program has had a significant impact on the likelihood of HipDVT (p for joint test of interaction term: 0.03), and KneeDVT (p for joint test of interaction term: <0.01). Importantly, the Program did not have a significant impact on the surgical any-or-none indicator.

The between-hospital variance (Random effects in the Model parameters section of Table 9) for the five primary outcomes ranged from 0.28 for the surgical any-or-none indicator to 0.78 for KneeDVT. This is proportional to the intraclass correlation coefficient (ICC) which represents the proportion of the total variance that is explained by hospital-level variability²⁵⁹. Importantly, despite a small intraclass correlation coefficient, the chi-

square statistic for the test of clustering is significant – thereby providing evidence that the between-hospital variance in primary outcomes is non-zero.

Appendix Table A 3 presents the interaction effects from the multivariate regression models for the secondary outcomes in each cohort. In the medical cohort, the Program was associated with a 70% decline in the predicted odds of UTI (COR: 0.30; 95% CI: 0.12 to 0.75; p-value for joint test: 0.09) among MUR quartile 3 hospitals as compared to MUR quartile 1 hospitals. In the surgical cohort, the Program was associated with an increase in the predicted odds of BSI among MUR quartile 3 (COR: 7.40; 95% CI: 1.23 to 44.55; p-value for joint test: 0.04) and quartile 4 (COR: 8.75; 95% CI: 1.74 to 44.03) hospitals.

Table 9: Multivariate odds ratios from hierarchical logistic regression models for the Program effects on the primary outcomes¹

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none</i> ³	<i>Any-or-none</i>	<i>SpineSSI</i> ⁴	<i>HipDVT</i> ⁵	<i>KneeDVT</i> ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Program⁶: Main effect					
Pre-	Ref	Ref	Ref	Ref	Ref
Post-	1.24 [0.83,1.86]	0.61 [0.30,1.21]	2.27 [0.90,5.73]	1.30 [0.62,2.77]	1.60 [0.67,3.82]
Medicare Utilization Ratio (MUR) Quartile⁷: Main effect					
Quartile 1: <=35.25%	Ref	Ref	Ref	Ref	Ref
Quartile 2: >35.25% & <=46.64%	1.55* [1.03,2.34]	1.09 [0.63,1.90]	2.38* [1.13,5.00]	1.98** [1.28,3.06]	1.83 [0.96,3.49]
Quartile 3: >46.64% & <=56.86%	1.03 [0.58,1.83]	0.55 [0.29,1.05]	1.44 [0.61,3.40]	1.84* [1.13,2.99]	2.48 [0.31,19.93]
Quartile 4: >56.86% & <=80.33%	1.23 [0.74,2.04]	0.61 [0.31,1.21]	1.85 [0.52,6.56]	1.82* [1.05,3.15]	1.91 [0.75,4.84]
Program x MUR: Interaction effects					
Post- x Quartile 2	0.57** [0.38,0.87]	1.07 [0.54,2.09]	0.34* [0.14,0.82]	0.56* [0.34,0.91]	0.71 [0.46,1.10]
Post- x Quartile 3	0.72 [0.45,1.17]	1.71 [0.80,3.64]	0.41* [0.18,0.94]	0.46** [0.27,0.79]	0.36** [0.18,0.74]
Post- x Quartile 4	0.69 [0.40,1.20]	1.38 [0.55,3.45]	0.41 [0.16,1.05]	0.51* [0.26,1.00]	0.70 [0.34,1.46]
p-value for joint test of interaction	0.07	0.40	0.07	0.03	0.00
Model parameters					
Inpatient stays	531,495	173,512	21,384	88,355	99,204
Hospitals	153	147	112	152	151
Random effects	0.41	0.28	0.26	0.34	0.78
Intraclass correlation coefficient	0.05	0.02	0.02	0.03	0.16
Likelihood ratio test					
Chi-square statistic	64.04	6.27	0.78	20.92	182.95
p-value	<0.001	0.01	0.19	<0.001	<0.001

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

Figure 6: Multivariate odds ratios from models evaluating the impact of the Hospital-Acquired Condition Program

3.4. Specific Aim 2.2

This aim evaluates the overall impact of the Program across hospitals with varying MUR and varying financial strength. The financial strength is represented by the operating profit margin (Hypothesis 2.2.1) and the DPP (Hypothesis 2.2.2).

3.4.1. Hypothesis 2.2.1

Controlling for relevant variables, the HAC Program is associated with a greater decline in HAC likelihood among Medicare beneficiaries admitted to NY hospitals that are at a higher risk of financial losses from the Program, and that this association is stronger among hospitals with higher profit margins.

The results of this analysis for the primary outcomes are tabulated in Table 10, and are discussed below. Appendix Table A 4 presents the results of this analysis for the secondary outcomes in each cohort.

Program main effects:

For MUR quartile 1 hospitals with negative profit margin, the Program was not associated with significant changes in the odds of HACs.

MUR main effects:

For negative profit margin hospitals in the pre-Program phase, the relative odds of the surgical any-or-none indicator were lower for MUR quartile 4 hospitals (COR: 0.19; 95% CI: 0.04 to 0.81) as compared to MUR quartile 1 hospitals. Stays in MUR quartile 2

hospitals had higher odds of SpineSSI (COR: 2.88; 95% CI: 1.19 to 6.98) and HipDVT (COR: 1.88; 95% CI: 1.08 to 3.27) as compared to MUR quartile 1 hospitals.

Operating profit margin main effects:

For MUR quartile 1 hospitals in the pre-Program phase, the odds of HACs were not significantly different between hospitals with negative and positive profit margins.

Program x MUR Quartile: 2-way interaction effects

For hospitals with negative profit margins, MUR quartile 2 hospitals (COR: 0.49; 95% CI: 0.26 to 0.92) had lower predicted odds of the medical any-or-none indicator in the post-Program phase as compared to MUR Quartile 1 hospitals.

Program x Operating profit margin: 2-way interaction effects

For MUR Quartile 1 hospitals, the Program was not associated with significant differences in the odds of HACs across hospitals with positive or negative profit margins.

MUR quartile x Operating profit margin: 2-way interaction effects

In the pre-Program phase, the Program was not associated with significant differences in the odds of HACs across hospitals with varying MUR and profit margins.

Program x MUR quartile x Operating profit margin: 3-way interaction effects

From multivariate analyses, the impact of the Program on HACs in hospitals with varying MUR load did not vary significantly between hospitals with positive and negative profit margins.

Table 10: Multivariate odds ratios for the primary outcomes¹ among hospitals with differential operating profit margins

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Program⁶: Main effect					
Pre-	Ref	Ref	Ref	Ref	Ref
Post-	1.13 [0.75,1.68]	0.64 [0.25,1.62]	2.65 [0.63,11.08]	1.03 [0.47,2.26]	1.85 [0.57,6.01]
Medicare Utilization Ratio (MUR) Quartile⁷: Main effect					
Quartile 1: <=35.25%	Ref	Ref	Ref	Ref	Ref
Quartile 2: >35.25% & <=46.64%	1.51 [0.93,2.47]	1.75 [0.81,3.78]	2.88* [1.19,6.98]	1.88* [1.08,3.27]	1.62 [0.73,3.59]
Quartile 3: >46.64% & <=56.86%	0.99 [0.52,1.90]	0.58 [0.21,1.61]	1.87 [0.64,5.47]	1.71 [0.93,3.17]	1.36 [0.32,5.69]
Quartile 4: >56.86% & <=80.33%	0.83 [0.39,1.79]	0.19* [0.04,0.81]	3.62 [0.83,15.88]	1.31 [0.61,2.81]	2.12 [0.62,7.27]
Operating profit margin: Main effect					
Negative	Ref	Ref	Ref	Ref	Ref
Positive	0.73 [0.31,1.73]	1.28 [0.47,3.48]	1.45 [0.55,3.86]	0.92 [0.58,1.47]	0.83 [0.37,1.87]
Program x MUR quartile: 2-way interaction					
Post- x Quartile 2	0.49* [0.26,0.92]	0.68 [0.26,1.79]	0.27 [0.06,1.12]	0.84 [0.43,1.62]	0.79 [0.38,1.64]
Post- x Quartile 3	0.74 [0.44,1.26]	2.08 [0.67,6.42]	0.48 [0.07,3.38]	0.55 [0.24,1.25]	0.88 [0.32,2.41]
Post- x Quartile 4	0.78 [0.38,1.61]	4.22 [0.74,23.93]	0.23 [0.02,2.84]	0.63 [0.17,2.27]	0.37 [0.12,1.14]
Program x Operating margin: 2-way interaction					
Post- # Positive margin	1.38 [0.53,3.57]	0.91 [0.25,3.31]	0.51 [0.09,2.81]	1.32 [0.63,2.76]	0.71 [0.28,1.80]
MUR quartile x Operating margin: 2-way interaction					
Quartile 2 # Positive margin	1.11 [0.44,2.83]	0.47 [0.15,1.48]	0.64 [0.17,2.39]	1.02 [0.57,1.79]	1.26 [0.50,3.15]
Quartile 3 # Positive margin	0.92 [0.30,2.80]	1.07 [0.28,4.06]	0.55 [0.13,2.32]	1.23 [0.57,2.66]	3.55 [0.99,12.79]
Quartile 2 # Positive margin	1.84 [0.58,5.80]	4.10 [0.74,22.61]	0.20 [0.04,1.08]	1.62 [0.73,3.59]	1.07 [0.29,3.94]
Program x MUR quartile x Operating margin: 3-way interaction					

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Post- # Quartile 2 # Positive margin	1.08 [0.36,3.27]	2.02 [0.47,8.73]	1.97 [0.25,15.51]	0.61 [0.25,1.49]	0.97 [0.33,2.82]
Post- # Quartile 3 # Positive margin	1.00 [0.32,3.19]	0.71 [0.14,3.47]	1.23 [0.13,12.06]	0.77 [0.27,2.25]	0.30 [0.08,1.19]
Post- # Quartile 4 # Positive margin	0.71 [0.21,2.40]	0.28 [0.03,2.45]	5.25 [0.27,102.23]	0.77 [0.19,3.06]	2.80 [0.57,13.69]
p-value for joint test of interaction	0.86	0.08	0.71	0.74	0.05
Model parameters					
Inpatient stays	529,307	171,516	21,024	87,491	98,034
Hospitals	152	146	112	152	150
Random effects	0.40	0.27	0.29	0.36	0.78
Intraclass correlation coefficient	0.05	0.02	0.03	0.04	0.16
Likelihood ratio test					
Chi-square test	57.64	5.43	1.22	23.35	181.90
p-value	<0.001	0.01	0.13	<0.001	<0.001

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

3.4.2. Hypothesis 2.2.2

Controlling for relevant variables, the HAC Program is associated with a greater decline in HAC likelihood among Medicare beneficiaries admitted to NY hospitals that are at a higher risk of financial losses from the Program, and that this association is weaker among hospitals with higher DPP.

The results of this analysis for the primary outcomes are tabulated in Table 11, and are discussed below. Appendix Table A 5 presents the results of this analysis for the secondary outcomes in each cohort.

Program main effects:

For MUR quartile 1 hospitals with DPP below the median, the Program was associated with an increase in the predicted odds of HipDVT (COR: 6.96; 95% CI: 2.63 to 18.43).

MUR main effects:

For hospitals with DPP below the median in the pre-Program phase, the relative odds of HipDVT were higher for MUR quartile 2 (COR: 4.44; 95% CI: 1.62 to 12.14), quartile 3 (COR: 4.41; 95% CI: 1.57 to 12.33), and quartile 4 (COR: 4.34; 95% CI: 1.51 to 12.46) hospitals.

DPP main effects:

For MUR quartile 1 hospitals in the pre-Program phase, the odds of the surgical any-or-none indicator (COR: 2.66; 95% CI: 1.18 to 6.02), and HipDVT (COR: 3.17; 95% CI: 1.23 to 8.14) were higher among hospitals with DPP above the median.

Program x MUR Quartile: 2-way interaction effects

For hospitals with DPP below the median, the odds of HipDVT were significantly lower among MUR quartile 2 (COR: 0.08; 95% CI: 0.04 to 0.18), quartile 3 (COR: 0.10; 95% CI: 0.04 to 0.23), and quartile 4 (COR: 0.10; 95% CI: 0.04 to 0.24) hospitals with the Program, as compared to MUR quartile 1 hospitals.

Program x DPP: 2-way interaction effects

For MUR Quartile 1 hospitals, the Program was associated with significant decline in the predicted odds of HipDVT (COR: 0.14; 95% CI: 0.06 to 0.31) among hospitals with DPP above the median.

MUR quartile x DPP: 2-way interaction effects

In the pre-Program phase, the predicted odds of the medical any-or-none indicator were significantly higher among MUR quartile 2 hospitals with DPP above median (COR: 2.79; 95% CI: 1.30 to 5.99), and the predicted odds of KneeDVT were significantly higher among MUR quartile 4 hospitals with DPP above median (COR: 7.61; 95% CI: 1.21 to 47.78).

Program x MUR quartile x DPP: 3-way interaction effects

From multivariate analyses, the predicted odds of HipDVT were significantly higher among hospitals with higher MUR load and above median DPP (For quartile 2 – COR: 10.51, 95% CI: 4.37 to 25.27; For quartile 3 – COR: 3.33, 95% CI: 1.02 to 10.83; p-value for joint test: <0.01) with the onset of the Program.

Table 11: Multivariate odds ratios for the primary outcomes¹ among hospitals with differential disproportionate patient percent

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Program⁶: Main effect					
Pre-	Ref	Ref	Ref	Ref	Ref
Post-	1.00 [0.50,2.01]	1.71 [0.56,5.19]	1.50 [0.28,8.08]	6.96*** [2.63,18.43]	0.85 [0.14,5.20]
Medicare Utilization Ratio (MUR) Quartile⁷: Main effect					
Quartile 1: <=35.25%	Ref	Ref	Ref	Ref	Ref
Quartile 2: >35.25% & <=46.64%	0.71 [0.32,1.56]	2.07 [0.79,5.42]	3.84 [0.73,20.30]	4.44** [1.62,12.14]	1.57 [0.44,5.52]
Quartile 3: >46.64% & <=56.86%	0.85 [0.39,1.82]	1.45 [0.57,3.71]	1.74 [0.37,8.23]	4.41** [1.57,12.33]	1.71 [0.11,25.69]
Quartile 4: >56.86% & <=80.33%	0.93 [0.45,1.96]	1.58 [0.65,3.83]	2.29 [0.42,12.54]	4.34** [1.51,12.46]	1.12 [0.25,5.12]
Disproportionate Patient Percent (DPP) Median: Main effect					
Below median	Ref	Ref	Ref	Ref	Ref
Above median	0.60 [0.31,1.16]	2.66* [1.18,6.02]	1.37 [0.28,6.77]	3.17* [1.23,8.14]	0.64 [0.16,2.46]
Program x MUR quartile: 2-way interaction					
Post- x Quartile 2	0.92 [0.41,2.06]	0.52 [0.16,1.73]	0.36 [0.07,1.71]	0.08*** [0.04,0.18]	1.32 [0.34,5.13]
Post- x Quartile 3	0.89 [0.46,1.73]	0.59 [0.18,1.98]	0.71 [0.15,3.41]	0.10*** [0.04,0.23]	0.67 [0.12,3.68]
Post- x Quartile 4	0.85 [0.39,1.87]	0.49 [0.15,1.61]	0.57 [0.11,2.96]	0.10*** [0.04,0.24]	1.28 [0.32,5.11]
Program x DPP median: 2-way interaction					
Post- # DPP above median	1.17 [0.59,2.30]	0.32 [0.09,1.14]	1.73 [0.40,7.57]	0.14*** [0.06,0.31]	2.07 [0.48,9.04]
MUR quartile x DPP median: 2-way interaction					

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Quartile 2 # DPP above median	2.79** [1.30,5.99]	0.56 [0.20,1.55]	0.49 [0.07,3.39]	0.37 [0.12,1.07]	1.07 [0.28,4.07]
Quartile 3 # DPP above median	1.07 [0.55,2.06]	0.30 [0.07,1.24]	1.70 [0.23,12.36]	0.48 [0.15,1.54]	1.04 [0.09,12.02]
Quartile 4 # DPP above median	1.33 [0.62,2.86]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.39 [0.29,6.56]	7.61* [1.21,47.78]
Program x MUR quartile x DPP median: 3-way interaction					
Post- # Quartile 2 # DPP above median	0.62 [0.27,1.44]	2.01 [0.48,8.32]	0.98 [0.16,6.10]	10.51*** [4.37,25.27]	0.47 [0.11,2.10]
Post- # Quartile 3 # DPP above median	1.00 [1.00,1.00]	3.81 [0.67,21.56]	1.00 [1.00,1.00]	3.33* [1.02,10.83]	0.52 [0.07,3.82]
Post- # Quartile 4 # DPP above median	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	4.38 [0.28,67.83]	0.32 [0.03,3.64]
p-value for joint test of interaction	0.27	0.32	0.98	0.00	0.72
Model parameters					
Inpatient stays	530,574	173,031	21,211	88,355	99,204
Hospitals	153	147	112	152	151
Random effects	0.42	0.28	0.25	0.35	0.81
Intraclass correlation coefficient	0.05	0.02	0.02	0.04	0.16
Likelihood ratio test					
Chi-square statistic	57.04	5.69	0.46	21.56	189.05
p-value	<0.001	0.01	0.25	<0.001	<0.001

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases

undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

3.5. Specific Aim 2.3

This aim evaluates the overall impact of the Program across hospitals with varying MUR and varying EHR capabilities, which is represented by the EMRAM stage for a hospital.

3.5.1. Hypothesis 2.3

Controlling for relevant variables, the HAC Program is associated with a greater decline in HAC likelihood among Medicare beneficiaries admitted to NY hospitals that are at a higher risk of financial losses from the Program, and that this association is stronger among hospitals with robust EHR capabilities.

The results of this analysis for the primary outcomes are tabulated in Table 12, and are discussed below. Appendix Table A 6 presents the results of this analysis for the secondary outcomes in each cohort.

Program main effect

For MUR quartile 1 hospitals with EMRAM Stage 0/1 EHRs, the Program was associated with a significant decrease in the predicted odds of surgical any-or-none indicator (COR: 0.13; 95% CI: 0.02 to 0.75).

MUR main effects:

For hospitals with EMRAM Stage 0/1 EHRs in the pre-Program phase, there were no significant differences in the predicted odds of HACs across hospitals with varying MUR load.

EMRAM main effects:

For MUR quartile 1 hospitals in the pre-Program phase, the odds of the surgical any-or-none indicator were higher among EMRAM stage 3+ hospitals (COR: 13.41; 95% CI: 1.43 to 125.48), and the odds of SpineSSI were lower for EMRAM Stage 2 hospitals (COR: 0.21; 95% CI: 0.09 to 0.49) as compared to EMRAM Stage 1 hospitals.

Program x MUR Quartile: 2-way interaction effects

For hospitals with EMRAM Stage 0/1 EHRs, the odds of SpineSSI were significantly lower among MUR quartile 3 (COR: 0.22; 95% CI: 0.06 to 0.85) and quartile 4 (COR: 0.32; 95% CI: 0.13 to 0.79) hospitals in the post-Program phase as compared to MUR quartile 1 hospitals.

Program x EMRAM Stage: 2-way interaction effects

For MUR Quartile 1 hospitals, the Program was associated with significantly higher predicted odds of the surgical any-or-none indicator among EMRAM Stage 2 hospitals (COR: 5.06; 95% CI: 1.02 to 25.04) as compared to EMRAM Stage 0/1 hospitals.

MUR quartile x EMRAM Stage: 2-way interaction effects

In the pre-Program phase, the odds of the surgical any-or-none indicator were significantly lower among MUR quartile 2 hospitals with EMRAM Stage 3 or higher EHRs (COR: 0.10; 95% CI: 0.01 to 0.99), among MUR quartile 3 hospitals with EMRAM Stage 3 or higher EHRs (COR: 0.07; 95% CI: 0.01 to 0.77), and among MUR quartile 4 hospitals with MUR Stage 2 EHRs (COR: 0.11; 95% CI: 0.01 to 1.00). In the pre-Program phase, the odds of HipDVT were significantly lower among MUR quartile 3

hospitals with EMRAM Stage 3+ EHRs (COR 0.32; 95% CI: 0.10 to 1.00), and among MUR quartile 4 hospitals with EMRAM Stage 3+ EHRs (COR: 0.28; 95% CI: 0.11 to 0.70). In the same phase, the odds of SpineSSI were significantly higher among MUR quartile 2 hospitals with EMRAM Stage 2 EHRs (COR: 4.90; 95% CI: 1.60 to 14.96).

Program x MUR quartile x EMRAM Stage: 3-way interaction effects

From multivariate analyses, the HAC Program was not associated with significant differences in the odds of HACs across hospitals with varying MUR load and EHR capabilities.

Table 12: Multivariate odds ratios for the primary outcomes¹ among hospitals with differential Electronic Health Record (EHR) capabilities

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none</i> ³	<i>Any-or-none</i>	<i>SpineSSI</i> ⁴	<i>HipDVT</i> ⁵	<i>KneeDVT</i> ⁶
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Program⁶: Main effect					
Pre-	Ref	Ref	Ref	Ref	Ref
Post-	0.84 [0.13,5.29]	0.13* [0.02,0.75]	1.77 [0.27,11.65]	1.42 [0.42,4.81]	2.06 [0.30,14.35]
Medicare Utilization Ratio (MUR) Quartile⁷: Main effect					
Q1: <=35.25%	Ref	Ref	Ref	Ref	Ref
Q2: >35.25% & <=46.64%	1.29 [0.26,6.54]	4.14 [0.51,33.62]	0.93 [0.38,2.24]	2.27 [0.85,6.10]	1.54 [0.07,34.32]
Q3: >46.64% & <=56.86%	0.77 [0.14,4.31]	2.13 [0.26,17.63]	0.60 [0.19,1.92]	1.81 [0.63,5.19]	4.54 [0.19,110.00]
Q4: >56.86% & <=80.33%	1.04 [0.20,5.48]	3.65 [0.44,30.10]	1.66 [0.42,6.58]	2.11 [0.77,5.75]	0.71 [0.10,4.90]
EMRAM Stage of EHR: Main effect					
Stage 0/1	Ref	Ref	Ref	Ref	Ref
Stage 2	0.75 [0.15,3.80]	2.70 [0.35,20.98]	0.21*** [0.09,0.49]	1.09 [0.45,2.66]	0.86 [0.07,10.61]
Stage 3+	1.11 [0.46,2.66]	13.41* [1.43,125.48]	0.29 [0.04,1.97]	0.81 [0.52,1.25]	0.56 [0.29,1.09]
Program x MUR quartile: 2-way interaction					
Post- # MUR Q2	0.88 [0.12,6.78]	2.59 [0.35,19.06]	0.44 [0.03,6.13]	1.59 [0.40,6.40]	0.25 [0.04,1.53]
Post- # MUR Q3	1.57 [0.24,10.34]	4.94 [0.68,35.77]	0.22* [0.06,0.85]	0.88 [0.20,3.85]	0.18 [0.01,5.75]
Post- # MUR Q4	0.30 [0.04,2.30]	1.37 [0.53,3.51]	0.32* [0.13,0.79]	1.17 [0.34,4.06]	0.79 [0.08,7.82]
Program x EMRAM Stage: 2-way interaction					
Post- # Stage 2	1.54 [0.22,11.01]	5.06* [1.02,25.04]	2.10 [0.50,8.90]	0.57 [0.19,1.77]	0.52 [0.08,3.46]
Post- # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
MUR quartile x EMRAM Stage: 2-way interaction					

Cohort²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none³</i>	<i>Any-or-none</i>	<i>SpineSSI⁴</i>	<i>HipDVT⁵</i>	<i>KneeDVT⁶</i>
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
MUR Q2 # Stage 2	1.05 [0.18,6.00]	0.21 [0.02,1.80]	4.90** [1.60,14.96]	0.83 [0.28,2.43]	1.33 [0.06,30.83]
MUR Q2 # Stage 3+	0.63 [0.20,2.01]	0.10* [0.01,0.99]	1.60 [0.08,34.20]	0.33 [0.10,1.11]	3.66 [0.98,13.70]
MUR Q3 # Stage 2	1.30 [0.20,8.34]	0.20 [0.02,1.79]	4.54 [0.97,21.28]	1.11 [0.30,4.07]	0.23 [0.01,8.75]
MUR Q3 # Stage 3+	0.60 [0.18,1.95]	0.07* [0.01,0.77]	4.52 [0.52,39.58]	0.32* [0.10,1.00]	0.88 [0.14,5.36]
MUR Q4 # Stage 2	1.11 [0.18,6.76]	0.11* [0.01,1.00]	1.13 [0.26,4.87]	0.81 [0.27,2.37]	4.14 [0.26,65.11]
MUR Q4 # Stage 3+	3.03 [0.66,13.84]	0.20 [0.02,1.89]	2.89 [0.47,17.60]	0.28** [0.11,0.70]	2.44 [0.46,12.85]
Program x MUR quartile x EMRAM Stage: 3-way interaction					
Post- # MUR Q2 # Stage 2	0.72 [0.08,6.68]	0.48 [0.07,3.55]	0.55 [0.04,8.62]	0.32 [0.07,1.56]	2.66 [0.45,15.78]
Post- # MUR Q2 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q3 # Stage 2	0.41 [0.05,3.37]	0.46 [0.07,3.04]	1.00 [1.00,1.00]	0.80 [0.11,5.94]	7.72 [0.23,256.87]
Post- # MUR Q3 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q4 # Stage 2	2.04 [0.21,19.52]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	0.61 [0.13,2.85]	0.70 [0.05,10.13]
Post- # MUR Q4 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
p-value for joint test of interaction	0.09	0.71	0.67	0.45	0.36
Model parameters					
Inpatient stays	529,224	171,659	21,196	87,626	98,180
Hospitals	152	146	110	151	149
Random effects	0.39	0.23	0.35	0.35	0.83
Intraclass correlation coefficient	0.04	0.02	0.04	0.04	0.17
Likelihood ratio test					
Chi-square statistic	53.32	3.03	2.21	22.02	189.77
p-value	<0.001	0.04	0.07	<0.001	<0.001

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group; EMRAM: Electronic Medical Record Adoption Model; EHR: Electronic Health Record

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

3.6. Stratified Analysis

To evaluate the impact of the Program on the risk of HACs for specific strata of patients, certain primary and secondary outcomes were evaluated in diagnosis- specific strata of the medical cohort. The interaction effects of this analysis are presented in Table 13.

With the onset of the Program, there was a significant decline in the odds of UTI among quartile 3 hospitals for the acute myocardial infarction stratum (COR: 0.23; 95% CI: 0.06 to 0.91). For the congestive heart failure stratum, there was a significant decline in the odds of the any-or-none indicator among quartile 3 hospitals with the onset of the Program (COR: 0.44; 95% CI: 0.22 to 0.88). For the pneumonia stratum, there was significant decline in the odds of the any-or-none indicator for quartile 2 hospitals (COR: 0.30; 95% CI: 0.11 to 0.83) and of the BSI indicator for quartile 2 (COR: 0.14; 95% CI: 0.04 to 0.47; p-value for joint test of interaction term: 0.01) and quartile 3 (COR: 0.28; 95% CI: 0.09 to 0.87; p-value for joint test of interaction term: 0.01) hospitals. For the stroke stratum, there was a significant decline in the odds of GLU for quartile 2 (COR: 0.17; 95% CI: 0.09 to 0.32; p-value for joint test of interaction term: <0.01) and 3 (COR: 0.27; 95% CI: 0.12 to 0.60; p-value for joint test of interaction term: <0.01) hospitals with the onset of the Program.

Table 13: Multivariate odds ratios for Hospital-Acquired Conditions in diagnosis-based strata of the medical cohort

Variables/Sub-Cohort	Any-or-none ²	BSI ³	UTI ⁴	FAL ⁵	GLU ⁶
Program x MUR¹: Interaction effects					
Acute myocardial infraction					
Post- x MUR Quartile 2	0.54 [0.25,1.19]	1.47 [0.51,4.24]	0.37 [0.08,1.75]	0.62 [0.20,1.97]	--
Post- x MUR Quartile 3	1.32 [0.45,3.85]	-- ⁷	0.23* [0.06,0.91]	0.71 [0.16,3.18]	--
Post- x MUR Quartile 4	1.15 [0.46,2.89]	2.27 [0.68,7.51]	0.72 [0.17,3.05]	1.05 [0.36,3.09]	--
p-value for joint test of interaction term	0.19	0.39	0.17	0.75	--
Congestive heart failure					
Post- x MUR Quartile 2	0.69 [0.37,1.26]	2.17 [0.82,5.78]	1.79 [0.75,4.23]	1.17 [0.52,2.65]	--
Post- x MUR Quartile 3	0.44* [0.22,0.88]	0.77 [0.23,2.52]	0.59 [0.21,1.62]	1.08 [0.42,2.81]	--
Post- x MUR Quartile 4	0.53 [0.27,1.04]	1.89 [0.50,7.18]	0.72 [0.18,2.94]	1.54 [0.64,3.73]	--
p-value for joint test of interaction term	0.09	0.25	0.03	0.74	--
Pneumonia					
Post- x MUR Quartile 2	0.30* [0.11,0.83]	0.14** [0.04,0.47]	--	0.87 [0.31,2.42]	--
Post- x MUR Quartile 3	0.52 [0.18,1.45]	0.28* [0.09,0.87]	--	1.22 [0.44,3.43]	--
Post- x MUR Quartile 4	0.73 [0.23,2.31]	0.89 [0.18,4.50]	--	1.71 [0.56,5.22]	--
p-value for joint test of interaction term	0.06	0.01	--	0.29	--
Stroke					
Post- x MUR Quartile 2	0.63 [0.24,1.62]	1.61 [0.56,4.60]	0.33 [0.05,2.29]	0.97 [0.38,2.48]	0.17*** [0.09,0.32]
Post- x MUR Quartile 3	0.87 [0.32,2.39]	3.14 [0.85,11.55]	0.35 [0.05,2.58]	2.24 [0.72,6.97]	0.27** [0.12,0.60]
Post- x MUR Quartile 4	0.41 [0.15,1.15]	1.00 [0.12,8.17]	0.44 [0.05,3.92]	0.8 [0.30,2.12]	--
p-value for joint test of interaction term	0.21	0.35	0.71	0.23	<0.01

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

¹ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

² Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility)

³ Catheter associated bloodstream infection ⁴ Catheter associated urinary tract infection ⁵ Falls and fractures ⁶ Poor glucose control

⁷ Failure to generate estimates due to non-convergence of the model

3.7. Sensitivity Analysis

3.7.1. Short- and long-term effects of the Program

The results for the sensitivity analysis examining the short-term and long-term effects of the Program are presented in Table 14. For reference, Section A of this table presents the interaction effects from the main analysis. Section B of the table presents the interaction effects for the short- (October 2008 to September 2010) and long-term (October 2010 to December 2012) analysis, along with the p-value for the joint test of the interaction term.

For the medical any-or-none indicator and HipDVT, the Program effects seen in the main analysis remain consistent in the short- and long-term phases. For the surgical any-or-none indicator, the Program is associated with an increase in the odds of the indicator for quartile 3 hospitals in the long-term. For SpineSSI, the overall intended Program effects that were noted in the main analysis are no longer significant when the post-Phase is divided into the short- and long-term. The Program effects for KneeDVT that were noted in the main analysis continue to remain consistent in the short-term but not in the long-term.

Table 14: Multivariate odds ratios for primary outcomes¹ in the short- and long-term post-Program phase

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Section A: Main Analysis – Program⁶ x MUR⁷: Interaction effects					
Post- x Quartile 2	0.57** [0.38,0.87]	1.07 [0.54,2.09]	0.34* [0.14,0.82]	0.56* [0.34,0.91]	0.71 [0.46,1.10]
Post- x Quartile 3	0.72 [0.45,1.17]	1.71 [0.80,3.64]	0.41* [0.18,0.94]	0.46** [0.27,0.79]	0.36** [0.18,0.74]
Post- x Quartile 4	0.69 [0.40,1.20]	1.38 [0.55,3.45]	0.41 [0.16,1.05]	0.51* [0.26,1.00]	0.70 [0.34,1.46]
p-value for joint test of interaction	0.07	0.40	0.07	0.03	0.00
Section B: Sensitivity Analysis – Program (Short- and long-term post-Program phases⁸) x MUR: Interaction effects					
Short term x Quartile 2	0.59* [0.37,0.95]	0.82 [0.40,1.65]	0.39 [0.15,1.00]	0.62 [0.35,1.11]	0.78 [0.42,1.44]
Short term x Quartile 3	0.70 [0.42,1.16]	1.33 [0.64,2.75]	0.42 [0.16,1.12]	0.47* [0.25,0.89]	0.43** [0.23,0.80]
Short term x Quartile 4	0.73 [0.41,1.30]	1.00 [0.40,2.46]	0.25 [0.06,1.04]	0.63 [0.30,1.33]	0.84 [0.35,2.03]
Long term x Quartile 2	0.55* [0.34,0.90]	1.75 [0.79,3.92]	0.33 [0.09,1.15]	0.48* [0.27,0.87]	0.65 [0.36,1.14]
Long term x Quartile 3	0.74 [0.42,1.31]	2.76* [1.07,7.08]	0.45 [0.13,1.53]	0.44** [0.24,0.80]	0.31 [0.09,1.02]
Long term x Quartile 4	0.65 [0.33,1.29]	2.43 [0.88,6.73]	0.46 [0.14,1.48]	0.39* [0.19,0.83]	0.57 [0.25,1.32]
p-value for joint test of interaction	0.28	0.03	0.25	0.08	0.05

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient

stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁸ Short-term phase extends from October 2008 to September 2010, and long-term phase extends from October 2010 to December 2012.

3.7.2. Redefining the Program phases

Table 15 presents findings from the analysis where the Program phases are redefined using certain important considerations. For reference, the Section A of the table provides the findings from the main analysis, Section B provides results where the post-Program phase is modelled to start in January 2009, and Section C presents results where observations with discharges in 2008 are excluded from the analysis.

For all the primary outcomes, the findings from the main analysis (Section A) are consistent with those from the sensitivity analysis where the post-phase is modelled to begin in January 2009 (Section B). For the sensitivity analysis that excludes observations with discharges in 2008 (Section C), the main analysis findings for HipDVT, KneeDVT, an surgical any-or-none indicator remain consistent, however the significant findings for the medical any-or-none indicator and SpineSSI that were noted in the main analysis are no longer significant in the sensitivity analysis.

Table 15: Multivariate odds ratios for primary outcomes¹ with redefined post-Program phases

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Section A: Main Analysis – Program⁶ x MUR⁷: Interaction effects					
Post- x Quartile 2	0.57** [0.38,0.87]	1.07 [0.54,2.09]	0.34* [0.14,0.82]	0.56* [0.34,0.91]	0.71 [0.46,1.10]
Post- x Quartile 3	0.72 [0.45,1.17]	1.71 [0.80,3.64]	0.41* [0.18,0.94]	0.46** [0.27,0.79]	0.36** [0.18,0.74]
Post- x Quartile 4	0.69 [0.40,1.20]	1.38 [0.55,3.45]	0.41 [0.16,1.05]	0.51* [0.26,1.00]	0.70 [0.34,1.46]
p-value for joint test of interaction	0.07	0.40	0.07	0.03	0.00
Section B: Sensitivity Analysis – Post-phase starts at 2009⁹ x MUR: Interaction effects					
Post- x Quartile 2	0.58* [0.39,0.88]	1.19 [0.58,2.41]	0.36* [0.13,0.98]	0.58* [0.36,0.96]	0.85 [0.56,1.29]
Post- x Quartile 3	0.61* [0.38,0.99]	1.49 [0.66,3.38]	0.41 [0.16,1.03]	0.50* [0.28,0.90]	0.43* [0.22,0.85]
Post- x Quartile 4	0.71 [0.42,1.21]	1.34 [0.57,3.17]	0.36 [0.11,1.13]	0.54 [0.29,1.03]	0.79 [0.39,1.59]
p-value for joint test of interaction	0.07	0.78	0.18	0.07	0.09
Section C: Sensitivity Analysis – Program (2008 excluded)¹⁰ x MUR: Interaction effects					
Post- x Quartile 2	0.61 [0.30,1.21]	0.96 [0.35,2.63]	--	0.45** [0.25,0.79]	0.59 [0.34,1.01]
Post- x Quartile 3	0.64 [0.28,1.44]	2.06 [0.64,6.66]	--	0.35** [0.17,0.72]	0.23** [0.09,0.61]
Post- x Quartile 4	1.14 [0.46,2.80]	5.23 [0.94,29.04]	0.35 [0.35,0.35]	0.36** [0.17,0.74]	0.58 [0.23,1.44]
p-value for joint test of interaction	0.25	0.08	1.00	0.01	0.03

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁸ Short-term phase extends from October 2008 to September 2010, and long-term phase extends from October 2010 to December 2012.

3.7.3. Redefining the MUR variable

For this group of sensitivity analyses, the MUR variable was redefined using quintile and tertile specifications. Table 16 contains the findings from this analysis. For reference, Section A presents the findings from the main analysis where the MUR variable is categorized into quartiles. Section B presents the analysis where the MUR is categorized into quintiles, and Section C presents the analysis for the MUR variable categorized into tertiles.

For SpineSSI, HipDVT, and KneeDVT (p-value for the joint test of interaction term: 0.04, <0.01, and <0.01 respectively), the findings from the quintile-specification analysis (Section B) are consistent with the findings from the main analysis (Section A). For the medical any-or-none indicator, the significant finding of the Program-associated decline is not observed in any of the categories in the quintile-specification. For the surgical any-or-none indicator, quintile 4 shows a significant increase in the odds as compared to quintile 1, in contrast to the findings from the main analysis where no significant change was observed among the MUR quartiles.

From the tertile-specification analysis (Section C), there is a significant decline in the predicted odds of HipDVT, and KneeDVT (p-value for the joint test of interaction term: 0.01 and 0.02 respectively) among tertile 2 hospitals with the introduction of the Program. However, the significant decline in the risk of the medical any-or-none indicator and SpineSSI that was noted in the main analysis is no longer found in this sensitivity analysis.

Table 16: Multivariate odds ratios for primary outcomes¹ with redefined Medicare Utilization Ratio variable

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Section A: Main Analysis – Program⁶ x MUR⁷: Interaction effects					
Post- x Quartile 2	0.57** [0.38,0.87]	1.07 [0.54,2.09]	0.34* [0.14,0.82]	0.56* [0.34,0.91]	0.71 [0.46,1.10]
Post- x Quartile 3	0.72 [0.45,1.17]	1.71 [0.80,3.64]	0.41* [0.18,0.94]	0.46** [0.27,0.79]	0.36** [0.18,0.74]
Post- x Quartile 4	0.69 [0.40,1.20]	1.38 [0.55,3.45]	0.41 [0.16,1.05]	0.51* [0.26,1.00]	0.70 [0.34,1.46]
p-value for joint test of interaction	0.07	0.40	0.07	0.03	0.00
Section B: Sensitivity Analysis – Program x MUR Quintile: Interaction effects					
Post- x Quintile 2	0.69 [0.42,1.14]	1.62 [0.80,3.28]	0.25** [0.10,0.63]	0.58* [0.34,0.99]	0.55 [0.29,1.07]
Post- x Quintile 3	0.74 [0.44,1.24]	1.79 [0.83,3.87]	0.32* [0.12,0.84]	0.34*** [0.20,0.57]	0.26*** [0.12,0.56]
Post- x Quintile 4	0.85 [0.48,1.49]	3.35** [1.40,7.99]	0.49 [0.14,1.66]	0.57 [0.26,1.24]	0.81 [0.43,1.54]
Post- x Quintile 5	0.70 [0.37,1.32]	1.68 [0.61,4.63]	0.28* [0.11,0.74]	0.39** [0.19,0.80]	0.61 [0.28,1.35]
p-value for joint test of interaction	0.64	0.11	0.04	<0.01	<0.01
Section C: Sensitivity Analysis – Program x MUR Tertile: Interaction effects					
Post- x Tertile 2	1.00 [0.68,1.46]	1.44 [0.82,2.51]	0.81 [0.36,1.82]	0.49** [0.32,0.76]	0.47** [0.28,0.81]
Post- x Tertile 3	1.04 [0.64,1.70]	1.59 [0.75,3.41]	0.92 [0.35,2.45]	0.62 [0.34,1.14]	1.06 [0.60,1.88]
p-value for joint test of interaction	0.98	0.36	0.87	0.01	0.02

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases

undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

3.7.4. Excluding low-volume hospitals

Table 17 presents the findings from the analysis where low volume hospitals are excluded from the analysis. For reference, section A of the table presents the findings from the main analysis. Section B of the table presents findings from the sensitivity analysis where hospitals with less than 20 observations for a year are excluded from the analysis. For all the primary outcomes with the exception of SpineSSI, the findings of the sensitivity analysis remain consistent with those from the main analysis. For SpineSSI, the significant decline in the rates that were seen among MUR quartile 2 and 3 hospitals in the main analysis are no longer significant in the sensitivity analysis.

Table 17: Multivariate odds ratios from analysis for primary outcomes¹ excluding hospitals with less than 20 observations a year

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Section A: Main Analysis – Program⁶ x MUR⁷: Interaction effects					
Post- x Quartile 2	0.57** [0.38,0.87]	1.07 [0.54,2.09]	0.34* [0.14,0.82]	0.56* [0.34,0.91]	0.71 [0.46,1.10]
Post- x Quartile 3	0.72 [0.45,1.17]	1.71 [0.80,3.64]	0.41* [0.18,0.94]	0.46** [0.27,0.79]	0.36** [0.18,0.74]
Post- x Quartile 4	0.69 [0.40,1.20]	1.38 [0.55,3.45]	0.41 [0.16,1.05]	0.51* [0.26,1.00]	0.70 [0.34,1.46]
p-value for joint test of interaction	0.07	0.40	0.07	0.03	0.00
Section B: Excluding hospitals with less than 20 observations a year					
Post- x Quartile 2	0.57** [0.38,0.87]	0.98 [0.49,1.99]	0.39 [0.15,1.05]	0.52* [0.31,0.88]	0.69 [0.45,1.06]
Post- x Quartile 3	0.73 [0.45,1.17]	1.50 [0.69,3.28]	0.68 [0.29,1.63]	0.45** [0.25,0.79]	0.36*** [0.23,0.56]
Post- x Quartile 4	0.69 [0.40,1.20]	1.25 [0.48,3.25]	0.47 [0.16,1.44]	0.44* [0.22,0.89]	0.71 [0.35,1.45]
p-value for joint test of interaction	0.07	0.55	0.23	0.03	<0.01

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient

stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

Chapter 4 - Discussion

The central aim of this study was to evaluate the overall impact of the HAC-POA Program on hospitals with varying Medicare load (Specific aim 2, hypothesis 1), and to determine if this impact varied across hospitals with varying financial strength (Specific aim 2, hypothesis 2) and digital capabilities (Specific aim 2, hypothesis 3). I used the conceptual model, theoretical framework and existing literature to develop the hypotheses. I hypothesized that the Program was associated with a decline in the likelihood of HACs and that in the post-Program phase hospitals at risk of higher financial losses from the Program would have lower likelihood of HACs as compared to their respective counterparts (Hypothesis 2.1). Furthermore, I hypothesized that in the post-Program phase, hospitals with positive operating margins (Hypothesis 2.2.1), lower disproportionate patient percent (Hypothesis 2.2.2), and advanced EHR capabilities (Hypothesis 2.3) would have lower risk of HACs as compared to their respective counterparts. Thus, the study aims to determine the role of financial disincentives in motivating hospitals to improve their care quality and reduce errors. The analysis was carried out separately in the medical and surgical cohorts that were defined using key diagnosis and procedure codes.

This section discusses the findings for the five primary outcomes i.e. the any-or-none indicators in each of the cohorts, surgical site infections among spine surgery patients (SpineSSI), and deep vein thrombosis or pulmonary embolism among hip and knee replacement patients (HipDVT and KneeDVT). A reference is made to key significant findings from secondary outcomes (BSI, UTI, FAL, GLU) (tables presented in the

Appendix) when appropriate. I begin by summarizing findings of the analysis, followed by a discussion on the meaning and potential explanations for the findings, and their policy implications. I end this section with a discussion on the limitations of the study.

4.1. Findings

A summary of the Program-associated findings are presented in Table 18 (only Program variables with significant effects are retained in the table), and are discussed in the following section.

4.1.1. Medical cohort

The any-or-none indicator in the medical cohort identifies the occurrence of any of the six HACs of interest, and is a composite measure for the safety performance of the hospital. The HAC-POA Program was significantly associated with a 43% decline in the relative odds of this indicator and a 70% decline in the odds of catheter-associated urinary tract infections among hospitals with higher Medicare load and consequently at higher risk for financial losses from the Program. Importantly, this group of hospitals had a higher baseline load of HACs. Importantly, hospital-level variability had a significant yet limited role to play in the risk of HACs, and patient factors explained a greater proportion of the variance in HACs.

The impact of the Program on the any-or-none indicator did not vary by the operating profit margin. However, for hospitals with negative operating profit margin, the MUR quartile 2 had a lower likelihood of the any-or-none indicator as compared to hospitals in MUR quartile 1 in the post-Program phase. This result was not significant for hospitals with positive operating margin. Furthermore, the impact of the Program on the any-or-none indicator did not vary by the disproportionate patient percent of the hospital, nor did it vary by the differing digital capabilities of a hospital.

From the stratified analysis, statistically significant decline in the likelihood of several HACs, with the exception of falls and trauma, was noted in the four diagnosis-based strata. From the sensitivity analyses, the impact of the Program on this indicator persisted both in the short- and the long-term; as well as when the program phases were redefined with the post-Program phase starting in January 2009. The effect also persisted when the models were re-estimated by excluding low-volume hospitals whose HAC rates may be outliers. However, the intended positive Program effects did not hold when the MUR variable were specified as quintiles and tertiles.

4.1.2. Surgical cohort

In the surgical cohort, the Program was associated with significant decline in relative odds of SpineSSI, (66% and 59% decline for MUR quartile 2 and 3 respectively), HipDVT (44%, 54% and 49% decline for MUR quartile 2 to 4 respectively), and KneeDVT (64% decline for MUR quartile 3 hospitals) among hospitals with higher Medicare patient load. Importantly, in the pre-Program phase hospitals with higher MUR had a higher likelihood of these HACs as compared to hospitals in MUR quartile 1.

The study did not identify significant Program effects that varied by the operating profit margin. Importantly, hospitals in MUR quartile 2 and 3 had significantly higher likelihood of HipDVT after Program implementation, and this effect was significant for hospitals with DPP above the median. Importantly, for hospitals with DPP below median, there was a significant decline in the odds of HipDVT with higher MUR load following Program implementation. The impact of the Program across hospitals with varying MUR load did not significantly vary across hospitals with differing digital capabilities. However, for

SpineSSI, among hospitals with low EHR capabilities, the Program was associated with decline in the likelihood of SpineSSI across hospitals with higher MUR.

From the sensitivity analyses, the significant impact of the Program on HipDVT and persisted both in the short- and the long-term; for KneeDVT it persisted in the short-term phase; and for SpineSSI significant effects were not noted when the Program phase was divided into short- and long-term. The findings of the analysis where the Program phases were redefined with the post-Program phase starting in January 2009, and where observations from 2008 were excluded yielded results that were fairly similar to those from the main analysis. The effects also persisted when the models were re-estimated by redefining MUR variable into tertile- and quintile-based categories, and by excluding low-volume hospitals whose HAC rates may be outliers (except for SpineSSI).

Table 18: Summary of findings for primary outcomes

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
(A) Main Analysis – Program⁶ x MUR⁷: Interaction effects.					
Hypothesis 2.1: Significant decline in HACs among higher MUR quartile hospitals					
Post- x Quartile 2	Significant - ⁸		Significant -	Significant -	
Post- x Quartile 3			Significant -	Significant -	Significant -
Post- x Quartile 4				Significant -	
(B) Main Analysis – Program x MUR x Operation margin					
Hypothesis 2.2.1: Impact of Program varies by operating margin of hospital					
Post- x Quartile 2	Significant -				
(C) Main Analysis – Program x MUR x Disproportionate patient percent					
Hypothesis 2.2.2: Impact of Program varies by disproportionate patient percent of hospital					
Post-				Significant -	
Post- x Quartile 2				Significant -	
Post- x Quartile 3				Significant -	
Post- x Quartile 4				Significant -	
Post- x DPP				Significant -	
Post- x Quartile 2 x DPP				Significant + ⁹	
Post- x Quartile 3 x DPP				Significant +	
(D) Main Analysis – Program x MUR x Digital capabilities					
Hypothesis 2.3: Impact of Program varies by digital capabilities of hospital					
Post-		Significant -			
Post- x Quartile 3			Significant -		
Post- x Quartile 4			Significant -		
Post- x EMRAM 2		Significant +			
(E) Stratified analysis – Acute myocardial infarction (AMI), Congestive Heart Failure (CHF), Pneumonia (PN), Stroke (STK)					

Cohort ²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
Outcome	Any-or-none ³	Any-or-none	SpineSSI ⁴	HipDVT ⁵	KneeDVT ⁵
Post- x Quartile 2	Significant – (PN)	Not applicable	Not applicable	Not applicable	Not applicable
Post- x Quartile 3	Significant – (CHF)				
(F) Sensitivity analysis – Short- and long-term phases					
Short-term x Quartile 2	Significant -				
Short-term x Quartile 3				Significant -	Significant -
Long-term x Quartile 2	Significant -			Significant -	
Long-term x Quartile 3		Significant +		Significant -	
Long-term x Quartile 4				Significant -	
(G) Sensitivity analysis – Post-phase starts in 2009					
Post- x Quartile 2	Significant -		Significant -	Significant -	
Post- x Quartile 3	Significant -			Significant -	Significant -
(H) Sensitivity analysis – Exclude 2008 observations					
Post- x Quartile 2				Significant -	
Post- x Quartile 3				Significant -	Significant -
Post- x Quartile 4				Significant -	
(I) Sensitivity analysis – MUR as quintile					
Post- x Quintile 2			Significant -	Significant -	
Post- x Quintile 3		Significant +	Significant -	Significant -	Significant -
Post- x Quintile 4					
Post- x Quintile 5			Significant -	Significant -	
(J) Sensitivity analysis – MUR as tertile					
Post- x Tertile 2				Significant -	Significant -
(K) Sensitivity analysis – Excluding low-volume hospitals					
Post- x Quartile 2	Significant -			Significant -	

Cohort²	Medical	Surgical	Surgical – Spine	Surgical – Hip Replacement	Surgical – Knee Replacement
<i>Outcome</i>	<i>Any-or-none³</i>	<i>Any-or-none</i>	<i>SpineSSI⁴</i>	<i>HipDVT⁵</i>	<i>KneeDVT⁵</i>
Post- x Quartile 3				Significant -	Significant -
Post- x Quartile 4				Significant -	

¹ Primary outcomes: Any-or-none indicator of the medical cohort (Column 2), Any-or-none indicator of the surgical cohort (Column 3), Surgical site infections among cases undergoing spine surgeries (Column 4), Deep vein thrombosis or pulmonary embolism among cases undergoing hip replacement (Column 5), Deep vein thrombosis or pulmonary embolism among cases undergoing knee replacement (Column 6).

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012. Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

³ Any-or-none indicator for the occurrence of six HACs (catheter associated bloodstream infection, catheter associated urinary tract infection, falls and fractures, poor glycemic control, air embolism, and blood incompatibility) in the respective cohort.

⁴ SpineSSI refers to the occurrence of surgical site infections among stays undergoing major spine surgeries of interest.

⁵ HipDVT and KneeDVT refer to the occurrence of deep vein thrombosis or pulmonary embolism among stays undergoing hip or knee replacement surgeries respectively.

⁶ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for the any-or-none indicators, and January 2005 to September 2008 for the Surgical site infections among cases undergoing Coronary Artery Bypass Graft or Spine Surgeries, and for Deep vein thrombosis or pulmonary embolism among cases under Hip or Knee Replacement surgeries. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁷ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁸ Implies significant decline in risk.

⁹ Implies significant increase in risk.

4.2. Discussion and Policy Implications

4.2.1. Importance of evaluating the HAC Program

Broadly, the findings of this study confirm that the Program was associated with significant decline in the likelihood of certain HACs, and that hospitals with higher Medicare load tend to be more responsive to the Program than hospitals with lower Medicare load.

The design of pay-for-performance ('carrots') or non-payment for non-performance ('stick') programs has been a subject of considerable discussion, without a clear consensus on which of these forms optimize results ²⁶⁰. Thus conducting a thorough evaluation of these Programs in a variety of settings and conditions will provide important insights for future policymaking. The design of the HAC-POA Program makes it unique in several ways. It is the first 'stick' program to target patient safety events in hospitals. While recent programs have incorporated a mix of bonuses and penalties in their design, the HAC-POA Program is the first program to exclusively assess disincentives for hospitals without awarding bonuses. Furthermore, while recent HAC-related legislations such as the HAC Reduction Program have directed disincentives towards poorly performing hospitals ²⁶¹, the HAC-POA Program is directed towards HACs in all eligible hospitals. Thus, rigorous empirical evaluation of this Program provides valuable insights for future reforms. The findings from this study provide evidence that supports the implementation of 'stick' programs to achieve safety targets.

4.2.2. Decline in the risk of HACs

The decline in the risk of certain HACs is an intended effect of the Program. There are several potential explanations for why this may have occurred. First, financial disincentives are a powerful motivator for hospitals to improve their quality and safety standards^{124, 262, 263}. As per the conceptual and theoretical models, hospitals are likely to have achieved these changes by changing their structures and processes, thereby resulting in better outcomes^{110, 248}. Second, the Program is credited with increasing the awareness of key hospital stakeholders towards HACs¹⁴⁶⁻¹⁴⁹. This is an important first step, because awareness of the hospital management about HACs and about their clinical and financial consequences is expected to translate into changes in structures and processes, which ultimately facilitate better outcomes²⁶⁴. Third, the availability of evidence-based guidelines and the standardization of preventive measures for UTI and BSI may have aided hospitals in updating their quality processes to prevent these infections from occurring¹⁴⁴. Importantly, our findings of decline in the likelihood of UTI and BSI are consistent with the findings from the study by Waters et al¹⁴⁴. Fourth, the HAC Program was initiated during a period of intense scrutiny on hospital quality and outcomes, especially infections. During this time, hospitals may have been aware about the financial and reputational implications of HACs – thereby motivating them to update their practices³⁴. We have controlled for other contemporaneous events thorough the study design.

Previous studies that have examined the Program have either found no evidence of the intended Program effects^{34, 142, 145} or they have found an intended decline in certain HACs such as deep vein thrombosis and pulmonary embolism¹⁴³. Importantly, none of the previous studies have evaluated the impact of the Program on falls and trauma, poor

glucose control or SpineSSI, and nor have they created a composite indicator for the overall HAC performance. This is the first study that addresses these areas of investigation, and the gaps in literature that have been previously discussed (Section 1.6.5).

These findings hold important directions for stakeholders in the quality improvement process. First, non-payment programs are associated with decline in the likelihood of HACs, and that this decline is seen among hospitals with higher Medicare patient load. Incorporating these findings into future reforms is likely to yield better outcomes. Second, given that contextual hospital effects have a significant yet limited role in explaining the variance in HACs, policy makers need to pay attention to the role of patient factors and within-hospital mechanisms that may facilitate or hinder desirable outcomes. It may help to qualitatively evaluate the positive changes that hospitals with improved HAC likelihood adopted with the introduction of the Program, and understand the feasibility of introducing such measures in other hospitals. For example, further evaluation of the potential for local programs to support sharing of quality improvement knowledge between hospitals, identification of quality improvement practices that can be emulated across hospitals, and the utility of state-level resources to strengthen hospital quality improvement efforts are some priority action areas that may result in gains in the value of hospital care. Third, concerted steps need to be taken to raise awareness about reforms and their implications. As gathered from previous qualitative studies, hospitals stakeholders were increasingly made aware about HACs and their implications through the HAC Program, and this may have played a role in the intended effects that we note. Fourth, key scientific associations need to accelerate the availability of evidence-based

guidelines not only for the prevention of HACs but also for the care of patients with key diagnoses and procedures. Adopting these guidelines is likely to provide greater returns. Fifth, this study examined four medical conditions and four surgical procedures for their responses to the Program. These form a small percentage of the admissions and surgeries for the inpatient Medicare population. While the HAC Program is applicable to most eligible inpatient admissions, designing 'sub-programs' that focus on other high impact conditions is likely to yield savings.

4.2.3. Absence of evidence for certain HACs

Despite falls and trauma being one of the most frequently occurring HACs, this study did not find evidence of a decline in its likelihood – neither in the main nor in the stratified analysis. The Program definition for falls and trauma is relatively broad, and includes fractures, dislocations, intracranial injuries, crushing injuries, and burns. The scope of this definition, the clinical diversity of the patients who are likely to encounter these adverse events, the range of preventive measures required to prevent these from occurring, and the financial trade-off in introducing such measures^{48, 49, 265-267} are some of the reasons why the study may not have identified significant Program effects for this HAC.

Considering the broad definition of this category of HACs, it is important to investigate the most commonly occurring conditions in this category, and especially conditions that are likely to yield to preventive strategies. By limiting the Program to such falls and trauma, hospitals are more likely to design more focused and effective preventive strategies.

4.2.4. Differential effects by operating margin

The financial health of a hospital is known to be an important indicator for quality outcomes^{109-111, 150}. Financially strong hospitals invest in structures and processes that facilitate quality outcomes. This study did not find evidence to support differential effects of the Program across hospitals with varying financial strength. However, since negative margin hospitals had a decline in the risk of the medical any-or-none indicator, it is likely that such hospitals invest in quality improvement efforts to avoid penalties that may further weaken their position. In contrast, since positive margin hospitals did not show a significant decline, it is likely that for these groups of hospitals, the losses from the Program were not significant enough to motivate further action, or that these groups of hospitals had little room for further improvement. Considering that hospitals with negative profit margins have taken steps to reduce HACs, supporting such hospitals financially through subsidies or by providing manpower and material support in the quality improvement process may prevent weakening of their financial conditions.

4.2.5. Differential effects by DPP (indicator for safety-net status)

The effect of the Program on the medical any-or-none indicator did not vary by the DPP or by the EHR capabilities of a hospital. This finding with respect to the DPP is contrary to common knowledge. Studies have shown that P4P programs tend to exacerbate the disparities in care between SN and non-SN hospitals^{149, 154, 157, 160, 165, 169-173}. However, most of these studies evaluated programs that offered bonuses and not penalties. Moreover, the differences in the study population under consideration make it challenging to match the findings between studies. Furthermore, since the clinical

conditions used to determine the cohort are already under considerable scrutiny by several pay-for-performance programs, it is likely that these groups are well positioned to prevent and address complications from care. Importantly, the HAC Program is unlikely to have exacerbated disparities in HAC outcomes, and that the safety-net status and digital capability of a hospital is unlikely to interact with the HAC Program to influence outcomes.

However, for HipDVT, the Program was associated with an increased risk among hospitals with DPP above the median. This finding is consistent with previous knowledge about payment reforms deepening the disparities between SN and non-SN hospitals. Hip replacements procedures are resource intensive procedures – both in terms of manpower and materials. Achieving high-quality outcomes following this surgery requires a coordinated systems approach, which in turn requires investments in quality. Patients admitted to safety-net hospitals are known to have severe clinical conditions, and these hospitals have limited financial resources. Thus, it is likely that the losses sustained from the HAC Program may have further constrained their resources resulting in poorer outcomes.

4.2.6. Differential effects by EHR capabilities

The study did not find evidence to support the differential Program effects across hospitals with varying digital capabilities. EHRs are tools that have been shown to support the provision of high-quality care ¹⁷⁴⁻¹⁸³. By standardizing the documentation of information, by facilitating access, by supporting the clinical decision-making process, EHRs are well positioned to provide critical information to providers in a timely manner.

Their role in influencing the likelihood of HACs is related to their ability to support the delivery of evidence-based care as a part of the prevention process. This study utilized the Electronic Medical Record Adoption Model (EMRAM) stage as an indicator to represent the use of EHRs. This indicator is generated from information provided by information technology officers or key administrators in a hospital. While an EHR may be implemented and in use in a hospital, the extent to which hospitals make use of the digital capabilities is debatable. Further research is needed to understand the translation of guidelines for preventing HACs into algorithms that can be implemented by EHRs.

4.2.7. Short- and long-term Program effects

Studies have shown that initial effects of pay-for-performance programs may taper off over time^{127, 129}. This may occur due to several reasons including that the incentives are no longer effective or that spillover effects into control groups obscures the identification of the program impact on the treatment group. Analyzing the long- and short-term effects of the HAC Program demonstrates that the disincentives of the Program were effective several years after Program implementation. Firstly, financial and reputation disincentives are powerful motivators for hospitals to undertake quality improvement measures. Importantly, improving HAC rates is a gradual process, with hospitals making modest improvements over a period of time. Thus, long-term sustained policy focus for reducing HACs are likely to yield benefits. Second, the start of the long-term phase also coincides with the implementation of the Affordable Care Act. It is likely that the focus on quality and safety metrics brought about by the Act may have further strengthened the response of the hospitals to the HAC Program.

4.2.8. Redefining the Program phases

Since the Program was introduced in October 2008, it is likely that hospitals may have initiated their practices in advance of the implementation or they may have deferred them to the start of 2009 to get a better understanding of the Program and its impact. Hence in the sensitivity analysis, models for the primary outcomes were re-estimated by excluding observations from 2008, and setting the post-Program phase to begin in 2009. Excluding observations from 2008 resulted in the Program effects becoming non-significant for the medical any-or-none indicator and for SpineSSI. This may be indicative of the fact that significant amount of decline may have occurred in advance of the Program being implemented ³⁴.

4.2.9. MUR specification

Several studies have used quartile-based specifications of key independent variables such as the disproportionate patient percent ²⁶⁸, and imaging scores ²⁶⁹ to study the differences in quality between high and low quartile hospitals. The quartile-based categorization of the MUR variable was pre-specified. The decision was based on the ability to determine and comment on the risk of HACs among hospitals with higher, intermediate, and lower load. The study identified significant decline in the HipDVT and KneeDVT in both the quintile and tertile specification, and in SpineSSI in the quintile specification. However the study did not identify significant decline in the medical any-or-none indicator in neither the quintile nor the tertile specification. These differences are explained by the change in the cut-offs for categorization of the MUR variable. Thus, inferences about significant changes may apply only to particular specifications of the MUR variable.

An alternate to categorization may have been using the continuous form of the MUR variable. By doing this we would the ratio to change by the same level among low and high MUR hospitals. However, since there are notable differences between these two groups of hospitals, they are likely to respond differently to the Program, and thus have differences in the change in risk.

4.2.10. Low volume hospitals

The volume-outcome relationship in hospitals is an important area of investigation ²⁷⁰. Hospitals that treat large number of patients with a particular diagnosis or procedure are more likely to have better outcomes on account of the practice and skill of those delivering care. Conversely, hospitals with low volume of patients are unlikely to have skilled resources – both manpower and material – such that the processes of care are standardized to achieve optimal outcomes. A reason for excluding these hospitals is that in low volume hospitals, the occurrence of even a single HAC may cause it to be an outlier, thereby influencing results. Re-estimation of models by excluding such low-volume hospitals did not change the interpretation of the estimates for most of the primary HACs. However, for SpineSSI, the findings from the main analysis were no longer significant indicating that SpineSSI in low volume hospitals may have played a role in the significant decline that was observed in the main analysis. This also supports the fact hospitals with low volume of spine surgeries may need to reconsider their quality programs, and seek quality-related support from institutions with high volume of spine surgeries.

4.3. Study limitations

Type of data:

This study utilizes administrative data, which is generated from billing records of inpatient stays. The variables used to identify HACs from this data are documented by medical coders based on the care that is provided and the information that is recorded in medical charts. Importantly, the primary purpose of this data is to optimize the billing revenue for each stay. While these datasets are accessible resources for undertaking studies with broad scope, they have certain noteworthy limitations. First, the ability of ICD-9 codes to effectively capture the clinical content of the charts may not be optimal. As an example, obese patients are at an increased risk of complications following hip, knee and spine surgeries. While the database contains an indicator of whether or not obesity was a comorbid condition, it does not provide details on the height, weight, and body mass index of the patient^{230, 271}, nor does it give an indication of the duration of obesity. Second, using ICD-9 codes to accurately document a condition is a combination of several key codes, and administrative datasets are known to have low sensitivity. As an example, coding for catheter-associated urinary tract infections requires the coder to code for both the use of the catheter and for the organism causing the infection. Failure to include one of these codes will result in a false negative entry²⁷². Third, administrative data lack variables that will enhance the risk adjustment methods. Examples of these include the vitals, physical examination findings, and laboratory values²⁷¹. Fourth, coding practices are dependent on the skills of the coder. Furthermore, it is well established that hospitals tend to code in ways that maximize their reimbursement²⁷³. Thus, these inter-hospitals variations in coding may further bias the findings. However, despite these limitations, the accessibility of the datasets makes it a reasonable

alternative to the medical record data. Importantly, CMS uses similar administrative datasets for identifying HACs and calculating financial disincentives.

Follow-up duration

The datasets that are used in this study capture the inpatient stays of patients. According to Program guidelines, this study identifies HACs based on their occurrence during the hospital stay. Other medical record datasets such as the National Surgical Quality Improvement Program (NSQIP) tracks the occurrence of complications for 30 days following surgery. It is likely that the symptoms of HACs may manifest several days after discharge from the hospital, and these may be treated in the outpatient or emergency setting. Furthermore, it is likely that patients may be readmitted for HACs that they acquired during their index stays. In such situations, the patient may be admitted with the hospital-acquired condition as a primary diagnosis that was present at admission. If this is to occur, the hospital is unlikely to be penalized for the HAC. Importantly, it is likely that our estimates of HACs are lesser than their actual likelihood. Despite this conservative estimate of HACs, the study has identified a significant impact of the Program.

Study design

The nested structure of the data supports the use of the multilevel study design. Because observations in a hospitals are likely to receive the same quality of care, and hence are likely to have similar outcomes, it is reasonable to state that there is a dependence among observations within hospitals. Failure to account for this dependence in the study design will result in underestimation of the standard errors, and

in the statistical significance of estimates when they are not actually significant. The multilevel study design can be implemented by using conditional methods such as fixed or random effects, and unconditional methods such as generalized estimating equations²⁵². Based on practical considerations (refer discussion in Section 2.4), random effects were used to account for variation within and between hospitals. However, effectively using random effects is contingent on meeting certain assumptions, key among which is that the unobserved confounders are not correlated with the covariates in the model. Failure to meet this assumption is likely to bias the estimates. As an example, the prevailing quality environment in a hospital is likely to influence the response of the hospital to the Program, and also on the HAC rates. Since a hospital that is more quality oriented is less likely to make changes in response to the Program (a likely negative correlation), and such a hospital is also likely to have higher quality and lower HACs (a likely negative correlation), the estimators are likely biased upwards / towards the null. An upward bias in this instance would mean that the estimated significant odds ratios are lesser than but closer to 1 than the true value. Thus, the estimates of the intended decline in HACs are likely conservative.

Causality

The central aim of a study evaluating policy effects would be to understand if the policy was solely responsible for the changes that occurred. Such inference would require an experimental setting where the magnitude of the 'dose' could be varied and the allocation of the 'dose' could be randomized. Since such a setup is unrealistic for evaluation of this Program, the study design used the difference-in-difference estimation routine. For studying the 'dose-response' of this Program, hospitals with lower Medicare

patient load were used as a proxy for hospitals that likely to face minimal disincentives from the Program. While there may have been other factors that have influenced the HAC rates, these are controlled using the difference-in-difference design and time fixed effects. While the quasi-experimental design using observational data does not confirm causality, it is difficult to identify other factors that may have led to this decline that are not controlled for in the analysis.

Geographic scope

This study examines the impact of the Program in New York State. The variation in the quality of care across states is a well-documented phenomenon. In addition to federal programs, the quality of care in a particular state is also the result of several local- and state-level initiatives. Importantly, NYS has had a rigorous focus on quality metrics through its report of the present-on-admission indicator, and its implementation of public reporting programs well in advance of national mandates. It is likely that the Program may not have had similar effects across other States, and thus the geographic generalizability of the findings is limited. Further research that examines the variation in the Program effects across states is needed to understand broad Program effects.

Structural and process measures

The dimensions of high quality care are optimal structure, processes, and outcomes²⁷⁴. Evaluating if the HAC-POA Program brought about improvements in quality and safety requires an evaluation of structures such as the human resource composition in a hospital, processes such as adoption of bundles for preventing hospital-acquired infections, and outcomes such as the occurrence of HACs. The focus of this study has

been on outcomes, which is also the focus of the Program. Since it does not examine the effects of the Program on hospital structures and processes, important advances in quality processes that hospitals may have made because of the Program are likely to be missed. However, there is reason to believe that the Program was associated with improvements in the quality environment and processes in hospitals. The existing qualitative peer-reviewed studies that surveyed infection preventionists and management in hospitals support the positive influence of the HAC Program.

4.4. Conclusion

The financial disincentives of the HAC-POA Program were associated with decline in the likelihood of certain HACs in New York hospitals. This decline was significant among hospitals at risk of higher financial losses from the Program. These policy effects were tested and hold in a variety of patient, hospital, and chronologic situations. Policy makers need to consider the strengths and drawbacks of this Program while formulating future payment reforms that target safety events in hospitals.

LIST OF REFERENCES

1. Blumenthal D. A Tale of Two Angels. *Ann Intern Med.* 2013.
2. Berwick DM, Hackbarth AD. Eliminating waste in US health care. *JAMA: the journal of the American Medical Association.* 2012;307(14):1513-1516.
3. Emanuel E, Tanden N, Altman S, et al. A systemic approach to containing health care spending. *N Engl J Med.* 2012;367(10):949-954.
4. Hoffman A EE. Reengineering US Health Care. *JAMA.* 2013;309(7):661-662. doi: 10.1001/jama.2012.214571.
5. McGlynn EA, Asch SM, Adams J, et al. The quality of health care delivered to adults in the United States. *N Engl J Med.* 2003;348(26):2635-2645.
6. Mechanic RE, Altman SH. Payment reform options: episode payment is a good place to start. *Health Aff (Millwood).* 2009;28(2):w262-71. doi: 10.1377/hlthaff.28.2.w262 [doi].
7. Rosenthal MB. Nonpayment for Performance? Medicare's New Reimbursement Rule. *New England Journal of Medicine.* 2007;357(16):1573-1575. doi: 10.1056/NEJMp078184.
8. Rosenthal MB, Fernandopulle R, Song HR, Landon B. Paying for quality: providers' incentives for quality improvement. *Health Aff.* 2004;23(2):127-141.
9. Rosenthal MB, Landon BE, Howitt K, Song HR, Epstein AM. Climbing up the pay-for-performance learning curve: where are the early adopters now?. *Health Aff (Millwood).* 2007;26(6):1674-1682. doi: 26/6/1674 [pii].
10. Damberg CL, Raube K, Teleki SS, Dela Cruz E. Taking stock of pay-for-performance: a candid assessment from the front lines. *Health Aff (Millwood).* 2009;28(2):517-525. doi: 10.1377/hlthaff.28.2.517 [doi].
11. Institute of Medicine. The healthcare imperative: lowering costs and improving outcomes: workshop series summary. National Academy Press; 2010.
12. Fineberg HV. A successful and sustainable health system—how to get there from here. *N Engl J Med.* 2012;366(11):1020-1027.
13. Kelley R. Where can \$700 billion in waste be cut annually from the US healthcare system?. Thomson Reuters.October. 2009.
14. PricewaterhouseCoopers' Health Research Institute. The Price of Excess: Identifying Waste in Healthcare Spending. <http://goo.gl/2Kj8uM>. Accessed 04/26, 2016.
15. Landrigan CP, Parry GJ, Bones CB, Hackbarth AD, Goldmann DA, Sharek PJ. Temporal trends in rates of patient harm resulting from medical care. *N Engl J Med.* 2010;363(22):2124-2134.
16. Leape LL, Berwick DM. Five years after to err is human. *JAMA: the journal of the American Medical Association.* 2005;293(19):2384-2390.
17. Pronovost PJ, Goeschel CA, Wachter RM. The wisdom and justice of not paying for “preventable complications”. *JAMA: The Journal of the American Medical Association.* 2008;299(18):2197-2199.

18. Titler MG. The Evidence for Evidence-Based Practice Implementation. In: Hughes RG, ed. Patient Safety and Quality: An Evidence-Based Handbook for Nurses. Rockville (MD): ; 2008.
19. Grover ML, Bracamonte JD, Kanodia AK, et al. Assessing adherence to evidence-based guidelines for the diagnosis and management of uncomplicated urinary tract infection. 2007;82(2):181-185.
20. Grol R. Successes and failures in the implementation of evidence-based guidelines for clinical practice. Med Care. 2001;39(8):II-46-II-54.
21. Ricart M, Lorente C, Diaz E, Kollef MH, Rello J. Nursing adherence with evidence-based guidelines for preventing ventilator-associated pneumonia. Crit Care Med. 2003;31(11):2693-2696.
22. Centers for Medicare and Medicaid Services. National Health Expenditure Fact Sheet. <https://goo.gl/Mse5EF>. Accessed 03/24, 2016.
23. Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL. The implications of regional variations in Medicare spending. Part 1: the content, quality, and accessibility of care. Ann Intern Med. 2003;138(4):273-287.
24. Leape LL, Brennan TA, Laird N, et al. The nature of adverse events in hospitalized patients: results of the Harvard Medical Practice Study II. N Engl J Med. 1991;324(6):377-384.
25. Thomas EJ, Lipsitz SR, Studdert DM, Brennan TA. The reliability of medical record review for estimating adverse event rates. Ann Intern Med. 2002;136(11):812-816.
26. Resnick NM, Marcantonio ER. How should clinical care of the aged differ?. The Lancet. 1997;350(9085):1157-1158.
27. Classen DC, Resar R, Griffin F, et al. 'Global trigger tool' shows that adverse events in hospitals may be ten times greater than previously measured. Health Aff. 2011;30(4):581-589.
28. Levinson DR, General I. Adverse events in hospitals: national incidence among Medicare beneficiaries. Department of Health & Human Services. 2010.
29. Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. N Engl J Med. 2009;360(14):1418-1428.
30. Hernandez AF, Greiner MA, Fonarow GC, et al. Relationship between early physician follow-up and 30-day readmission among Medicare beneficiaries hospitalized for heart failure. JAMA: The Journal of the American Medical Association. 2010;303(17):1716-1722.
31. Kohn LT, Corrigan JM, Donaldson MS. To err is human: building a safer health system. Vol 627. National Academies Press; 2000.
32. National Quality Forum. Serious Reportable Events in Healthcare - 2011 Update: A Consensus Report. http://www.qualityforum.org/Publications/2011/12/Serious_Reportable_Events_in_Health_care_2011.aspx. Accessed 09/05, 2013.

33. The Commonwealth Fund. Minnesota: Comprehensive Quality Improvement and Payment Reform. <http://goo.gl/gUbEyt>. Accessed 04/26, 2016.
34. Lee GM, Kleinman K, Soumerai SB, et al. Effect of Nonpayment for Preventable Infections in US Hospitals. *New England Journal of Medicine*. 2012;367(15):1428-1437.
35. Jha AK. Time to get serious about pay for performance. *JAMA*. 2013;309(4):347-348.
36. Rosenthal MB. Beyond pay for performance - Emerging models of provider-payment reform. *N Engl J Med*. 2008;359(12):1197-1200. doi: 10.1056/NEJMp0804658.
37. Wald HL, Kramer AM. Nonpayment for harms resulting from medical care - Catheter-associated urinary tract infections. *Jama-Journal of the American Medical Association*. 2007;298(23):2782-2784. doi: 10.1001/jama.298.23.2782.
38. Eappen S, Lane BH, Rosenberg B, et al. Relationship Between Occurrence of Surgical Complications and Hospital Finances. *JAMA*. 2013;309(15):1599-1606.
39. Leatherman S, Berwick D, Iles D, et al. The business case for quality: case studies and an analysis. *Health Aff*. 2003;22(2):17-30.
40. Centers for Medicare and Medicaid Services. 78 FR 50495 (Document Number: 2013-18956) August 19, 2013. <https://federalregister.gov/a/2013-18956>. Accessed 08/28, 2013.
41. Stone PW, Glied SA, McNair PD, et al. CMS Changes in Reimbursement for HAIs: Setting A Research Agenda. *Medical Care*. 2010;48(5):433-439. doi: 10.1097/MLR.0b013e3181d5fb3f.
42. Department of Health and Human Services. 77 FR 53257 (Document Number: 2010-19092; August 31, 2012). <https://goo.gl/NJOnSO>. Accessed 04/26, 2016.
43. Department of Health and Human Services. 73 FR 48434 (Document Number: E8-17914; August 19, 2008). <https://federalregister.gov/a/E8-17914>. Accessed 03/21, 2013.
44. QualityNet. Hospital-Acquired Conditions. <http://www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic%2FPage%2FQnetTier3&cid=12287594797672013>.
45. QualityNet. Specifications Manual for Hospital Inpatient Quality Reporting Program (Version 4.1). <http://goo.gl/vOPbKp>. Accessed 11/11, 2013.
46. RTI International. HAC-POA Evaluation: RTI Evaluation of FY 2011 Data. <http://goo.gl/IYWQhF>. Accessed 04/26, 2016.
47. Kandilov AM, Coomer NM, Dalton K. The Impact of Hospital-Acquired Conditions on Medicare Program Payments. *Medicare & Medicaid Research Review*. 2014;4(4).
48. Horwitz JR. Making profits and providing care: comparing nonprofit, for-profit, and government hospitals. *Health Aff (Millwood)*. 2005;24(3):790-801. doi: 24/3/790 [pii].
49. Chang T, Jacobson M. What is the mission of a not-for-profit hospital? Evidence from California's seismic retrofit mandate. Mimeo. 2008.
50. Weisbrod BA. The nonprofit economy. Harvard University Press; 2009.

51. Dranove D. Pricing by non-profit institutions: the case of hospital cost-shifting. *J Health Econ.* 1988;7(1):47-57.
52. Sloan FA, Vraciu RA. Investor-owned and not-for-profit hospitals: addressing some issues. *Health Aff.* 1983;2(1):25-37.
53. Dudley RA, Frolich A, Robinowitz DL, Talavera JA, Broadhead P, Luft HS. Strategies To Support Quality-based Purchasing: A Review of the Evidence. 2004.
54. Zhan C, Friedman B, Mosso A, Pronovost P. Medicare payment for selected adverse events: building the business case for investing in patient safety. *Health Aff.* 2006;25(5):1386-1393.
55. Hibbard JH. What can we say about the impact of public reporting? Inconsistent execution yields variable results. *Ann Intern Med.* 2008;148(2):160-161.
56. Hibbard JH, Stockard J, Tusler M. Hospital performance reports: impact on quality, market share, and reputation. *Health Aff.* 2005;24(4):1150-1160.
57. Mukamel DB, Mushlin AI. Quality of care information makes a difference: an analysis of market share and price changes after publication of the New York State Cardiac Surgery Mortality Reports. *Med Care.* 1998;36(7):945-954.
58. Romano PS, Marcin JP, Dai JJ, et al. Impact of public reporting of coronary artery bypass graft surgery performance data on market share, mortality, and patient selection. *Med Care.* 2011;49(12):1118-1125. doi: 10.1097/MLR.0b013e3182358c78 [doi].
59. Romano PS, Zhou H. Do well-publicized risk-adjusted outcomes reports affect hospital volume?. *Med Care.* 2004;42(4):367-377. doi: 00005650-200404000-00010 [pii].
60. Chaganti R, Sambharya R. Strategic orientation and characteristics of upper management. *Strategic Manage J.* 1987;8(4):393-401.
61. Hambrick DC, Mason PA. Upper echelons: The organization as a reflection of its top managers. *Academy of management review.* 1984;9(2):193-206.
62. Donabedian A. Evaluating the quality of medical care. *Milbank Mem Fund Q.* 1966;44(3):166-206.
63. Kaplan HC, Brady PW, Dritz MC, et al. The influence of context on quality improvement success in health care: a systematic review of the literature. *Milbank Q.* 2010;88(4):500-559.
64. Rosenthal MB, Frank RG. What is the empirical basis for paying for quality in health care?. *Medical Care Research and Review.* 2006;63(2):135-157.
65. Centers for Medicare and Medicaid Services. Acute Care Hospital Inpatient Prospective Payment System. <https://goo.gl/3DhmzC>. Accessed 02/10, 2016.
66. Brennan TA, Leape LL, Laird NM, et al. Incidence of adverse events and negligence in hospitalized patients: results of the Harvard Medical Practice Study I. *N Engl J Med.* 1991;324(6):370-376.
67. Brennan TA, Hebert LE, Laird NM, et al. Hospital characteristics associated with adverse events and substandard care. *JAMA: the journal of the American Medical Association.* 1991;265(24):3265-3269.

68. Keeler EB, Rubenstein LV, Kahn KL, et al. Hospital characteristics and quality of care. *JAMA: the journal of the American Medical Association*. 1992;268(13):1709-1714.
69. Hartz AJ, Krakauer H, Kuhn EM, et al. Hospital characteristics and mortality rates. *N Engl J Med*. 1989;321(25):1720-1725.
70. Tsilimingras D, Rosen AK, Berlowitz DR. Review Article: Patient Safety in Geriatrics: A Call for Action. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2003;58(9):M813-M819.
71. Lefevre F, Feinglass J, Potts S, et al. Iatrogenic complications in high-risk, elderly patients. *Arch Intern Med*. 1992;152(10):2074.
72. Thomas EJ, Brennan TA. Incidence and types of preventable adverse events in elderly patients: population based review of medical records. *BMJ*. 2000;320(7237):741-744.
73. Gawande AA, Thomas EJ, Zinner MJ, Brennan TA. The incidence and nature of surgical adverse events in Colorado and Utah in 1992. *Surgery*. 1999;126(1):66-75.
74. Gillick MR, Serrell NA, Gillick LS. Adverse consequences of hospitalization in the elderly. *Soc Sci Med*. 1982;16(10):1033-1038.
75. Silber JH, Williams SV, Krakauer H, Schwartz JS. Hospital and patient characteristics associated with death after surgery: a study of adverse occurrence and failure to rescue. *Med Care*. 1992;615-629.
76. Russo C, Steiner C, Spector W. Hospitalizations related to pressure ulcers among adults 18 years and older, 2006. *Healthcare Cost Utilization Project*. December. 2008.
77. Oliver D, Daly F, Martin FC, McMurdo ME. Risk factors and risk assessment tools for falls in hospital in-patients: a systematic review. *Age Ageing*. 2004;33(2):122-130.
78. Cohen B, Yoon Jeong Choi RN, MSN, Sandra Hyman R, E Yoko Furuya MD, MS, Elaine Larson R. Gender Differences in Risk of Bloodstream and Surgical Site Infections. *Journal of general internal medicine*. 2013;1-8.
79. Garibaldi RA, Burke JP, Dickman ML, Smith CB. Factors predisposing to bacteriuria during indwelling urethral catheterization. *N Engl J Med*. 1974;291(5):215-219.
80. Stamm WE. Catheter-associated urinary tract infections: epidemiology, pathogenesis, and prevention. *Am J Med*. 1991;91(3):S65-S71.
81. Metersky ML, Hunt DR, Kliman R, et al. Racial disparities in the frequency of patient safety events: results from the national medicare patient safety monitoring system. *Med Care*. 2011;49(5):504-510.
82. Fiscella K, Franks P, Meldrum S, Barnett S. Racial disparity in surgical complications in New York State. *Ann Surg*. 2005;242(2):151.
83. Hasan O, Orav EJ, Hicks LS. Insurance status and hospital care for myocardial infarction, stroke, and pneumonia. *Journal of Hospital Medicine*. 2010;5(8):452-459.
84. Lemaire A, Cook C, Tackett S, Mendes DM, Shortell CK. The impact of race and insurance type on the outcome of endovascular abdominal aortic aneurysm (AAA) repair. *Journal of vascular surgery*. 2008;47(6):1172-1180.

85. Weissman JS, Gatsonis C, Epstein AM. Rates of avoidable hospitalization by insurance status in Massachusetts and Maryland. *JAMA*. 1992;268(17):2388-2394.
86. Weissman JS, Vogeli C, Levy DE. The Quality of Hospital Care for Medicaid and Private Pay Patients. *Med Care*. 2013;51(5):389-395. doi: 10.1097/MLR.0b013e31827fef95.
87. Spencer CS, Gaskin DJ, Roberts ET. The quality of care delivered to patients within the same hospital varies by insurance type. *Health Aff (Millwood)*. 2013;32(10):1731-1739. doi: 10.1377/hlthaff.2012.1400 [doi].
88. Dimick JB, Pronovost PJ, Cowan Jr JA, Lipsett PA, Stanley JC, Upchurch Jr GR. Variation in postoperative complication rates after high-risk surgery in the United States. *Surgery*. 2003;134(4):534-540.
89. Dimick JB, Pronovost PJ, Cowan JA, Lipsett PA. Surgical volume and quality of care for esophageal resection: do high-volume hospitals have fewer complications?. *Ann Thorac Surg*. 2003;75(2):337-341.
90. Dang D, Johantgen ME, Pronovost PJ, Jenckes MW, Bass EB. Postoperative complications: does intensive care unit staff nursing make a difference?. *Heart & Lung: The Journal of Acute and Critical Care*. 2002;31(3):219-228.
91. Iezzoni LI, Daley J, Heeren T, et al. Identifying complications of care using administrative data. *Med Care*. 1994;700-715.
92. Mark BA, Harless DW, McCue M, Xu Y. A longitudinal examination of hospital registered nurse staffing and quality of care. *Health Serv Res*. 2004;39(2):279-300.
93. Escarce JJ, Kelley MA. Admission source to the medical intensive care unit predicts hospital death independent of APACHE II score. *JAMA*. 1990;264(18):2389-2394.
94. Naessens JM, Campbell CR, Shah N, et al. Effect of illness severity and comorbidity on patient safety and adverse events. *American Journal of Medical Quality*. 2012;27(1):48-57.
95. Rosen AK, Ash AS, McNiff KJ, Moskowitz MA. The importance of severity of illness adjustment in predicting adverse outcomes in the Medicare population. *J Clin Epidemiol*. 1995;48(5):631-643.
96. Rothschild JM, Bates DW, Leape LL. Preventable medical injuries in older patients. *Arch Intern Med*. 2000;160(18):2717.
97. Ghaferi AA, Birkmeyer JD, Dimick JB. Variation in hospital mortality associated with inpatient surgery. *N Engl J Med*. 2009;361(14):1368-1375.
98. Merkow RP, Ju MH, Chung JW, et al. Underlying reasons associated with hospital readmission following surgery in the United States. *JAMA*. 2015;313(5):483-495.
99. Barnett MJ, Kaboli PJ, Sirio CA, Rosenthal GE. Day of the week of intensive care admission and patient outcomes: a multisite regional evaluation. *Med Care*. 2002;40(6):530-539.
100. Kostis WJ, Demissie K, Marcella SW, Shao Y, Wilson AC, Moreyra AE. Weekend versus weekday admission and mortality from myocardial infarction. *N Engl J Med*. 2007;356(11):1099-1109.

101. Bell CM, Redelmeier DA. Mortality among patients admitted to hospitals on weekends as compared with weekdays. *N Engl J Med*. 2001;345(9):663-668.
102. Kind AJ, Jencks S, Brock J, et al. Neighborhood socioeconomic disadvantage and 30-day rehospitalization: a retrospective cohort study. *Ann Intern Med*. 2014;161(11):765-774.
103. Thiemann DR, Coresh J, Oetgen WJ, Powe NR. The association between hospital volume and survival after acute myocardial infarction in elderly patients. *N Engl J Med*. 1999;340(21):1640-1648.
104. Sheikh K, Bullock C. Urban-rural differences in the quality of care for Medicare patients with acute myocardial infarction. *Arch Intern Med*. 2001;161(5):737.
105. Eggleston K, Shen Y, Lau J, Schmid CH, Chan J. Hospital ownership and quality of care: what explains the different results in the literature?. *Health Econ*. 2008;17(12):1345-1362.
106. Cutler DM. The changing hospital industry: Comparing not-for-profit and for-profit institutions. University of Chicago Press; 2007.
107. Thomas EJ, Orav EJ, Brennan TA. Hospital ownership and preventable adverse events. *Journal of general internal medicine*. 2000;15(4):211-219.
108. Lipworth L, Lee J, Morris J. Case-fatality in teaching and non-teaching hospitals, 1956-1959. *The Challenge of Epidemiology: Issues and Selected Readings*. 2004;1(1):903-908.
109. Burstin HR, Lipsitz SR, Udvarhelyi IS, Brennan TA. The effect of hospital financial characteristics on quality of care. *JAMA: The Journal of the American Medical Association*. 1993;270(7):845-849.
110. Bazzoli GJ, Chen H, Zhao M, Lindrooth RC. Hospital financial condition and the quality of patient care. *Health Econ*. 2008;17(8):977-995.
111. Bazzoli GJ, Clement JP, Lindrooth RC, et al. Hospital financial condition and operational decisions related to the quality of hospital care. *Medical Care Research and Review*. 2007;64(2):148-168.
112. Rosenthal GE, Harper DL, Quinn LM, Cooper GS. Severity-adjusted mortality and length of stay in teaching and nonteaching hospitals. *JAMA: the journal of the American Medical Association*. 1997;278(6):485-490.
113. Ayanian JZ, Weissman JS. Teaching hospitals and quality of care: a review of the literature. *Milbank Q*. 2002;80(3):569-593.
114. Kessler DP, Geppert JJ. The effects of competition on variation in the quality and cost of medical care. *Journal of Economics & Management Strategy*. 2005;14(3):575-589.
115. Zwanziger J, Melnick GA. The effects of hospital competition and the Medicare PPS program on hospital cost behavior in California. *J Health Econ*. 1988;7(4):301-320.
116. Propper C, Burgess S, Green K. Does competition between hospitals improve the quality of care?: Hospital death rates and the NHS internal market. *Journal of Public Economics*. 2004;88(7):1247-1272.

117. Gowrisankaran G, Town RJ. Competition, Payers, and Hospital Quality¹. *Health Serv Res.* 2003;38(6p1):1403-1422.
118. The Commonwealth Fund. The Commonwealth Fund Health Care Opinion Leaders Survey: Assessing Health Care Experts' Views on Health Care Costs. <http://goo.gl/uZr75f>. Accessed 04/26, 2016.
119. Committee on Redesigning Health Insurance Performance Measures, Payment, and Performance Improvement Programs. *Performance Measurement: Accelerating Improvement (Pathways to Quality Health Care Series)*. The National Academies Press; 2006.
120. Grol RP, Bosch MC, Hulscher ME, Eccles MP, Wensing M. Planning and studying improvement in patient care: the use of theoretical perspectives. *Milbank Q.* 2007;85(1):93-138.
121. Epstein AM. Will pay for performance improve quality of care? The answer is in the details. *N Engl J Med.* 2012;367(19):1852-1853.
122. Van Herck P, De Smedt D, Annemans L, Remmen R, Rosenthal MB, Sermeus W. Systematic review: Effects, design choices, and context of pay-for-performance in health care. *Bmc Health Services Research.* 2010;10:247. doi: 10.1186/1472-6963-10-247.
123. Lindenauer PK, Remus D, Roman S, et al. Public reporting and pay for performance in hospital quality improvement. *N Engl J Med.* 2007;356(5):486-496. doi: 10.1056/NEJMsa064964.
124. Werner RM, Kolstad JT, Stuart EA, Polsky D. The Effect Of Pay-For-Performance In Hospitals: Lessons For Quality Improvement. *Health Aff.* 2011;30(4):690-698. doi: 10.1377/hlthaff.2010.1277.
125. Glickman SW, Ou F, DeLong ER, et al. Pay for performance, quality of care, and outcomes in acute myocardial infarction. *Jama-Journal of the American Medical Association.* 2007;297(21):2373-2380. doi: 10.1001/jama.297.21.2373.
126. Ryan AM. Effects of the Premier Hospital Quality Incentive Demonstration on Medicare Patient Mortality and Cost. *Health Serv Res.* 2009;44(3):821-842. doi: 10.1111/j.1475-6773.2009.00956.x.
127. Jha AK, Joynt KE, Orav EJ, Epstein AM. The Long-Term Effect of Premier Pay for Performance on Patient Outcomes. *N Engl J Med.* 2012;366(17):1606-1615. doi: 10.1056/NEJMsa1112351.
128. Sutton M, Nikolova S, Boaden R, Lester H, McDonald R, Roland M. Reduced mortality with hospital pay for performance in England. *N Engl J Med.* 2012;367(19):1821-1828.
129. Kristensen SR, Meacock R, Turner AJ, et al. Long-term effect of hospital pay for performance on mortality in England. *N Engl J Med.* 2014;371(6):540-548.
130. Hartmann CW, Hoff T, Palmer JA, Wroe P, Dutta-Linn MM, Lee G. The Medicare Policy of Payment Adjustment for Health Care-Associated Infections: Perspectives on Potential Unintended Consequences. *Medical Care Research and Review.* 2012;69(1):45-61. doi: 10.1177/1077558711413606.

131. Christianson JB, Leatherman S, Sutherland K. Lessons from evaluations of purchaser pay-for-performance programs a review of the evidence. *Medical Care Research and Review*. 2008;65(6 suppl):5S-35S.
132. Kavanagh KT, Cimiotti JP, Abusalem S, Coty M. Moving Healthcare Quality Forward With Nursing-Sensitive Value-Based Purchasing. *Journal of Nursing Scholarship*. 2012;44(4):385-395.
133. Rosenthal MB, Frank RG, Li Z, Epstein AM. Early experience with pay-for-performance. *JAMA: the journal of the American Medical Association*. 2005;294(14):1788-1793.
134. Werner RM, Goldman LE, Dudley RA. Comparison of change in quality of care between safety-net and non-safety-net hospitals. *Jama-Journal of the American Medical Association*. 2008;299(18):2180-2187. doi: 10.1001/jama.299.18.2180.
135. Agency for Healthcare Research and Quality. Never Events. <http://psnet.ahrq.gov/primer.aspx?primerID=3>. Accessed 11/19, 2013.
136. Berenson RA, Paulus RA, Kalman NS. Medicare's readmissions-reduction program—a positive alternative. *N Engl J Med*. 2012;366(15):1364-1366.
137. McNair PD, Luft HS, Bindman AB. Medicare's Policy Not To Pay For Treating Hospital-Acquired Conditions: The Impact. *Health Aff*. 2009;28(5):1485-1493. doi: 10.1377/hlthaff.28.5.1485.
138. McNutt R, Johnson TJ, Odwazny R, et al. Change in MS-DRG assignment and hospital reimbursement as a result of Centers for Medicare & Medicaid changes in payment for hospital-acquired conditions: is it coding or quality?. *Qual Manag Health Care*. 2010;19(1):17-24.
139. McHugh M, Martin TC, Orwat J, Van Dyke K. Medicare's policy to limit payment for hospital-acquired conditions: the impact on safety net providers. *J Health Care Poor Underserved*. 2011;22(2):638-647.
140. Stone PW, Pogorzelska M, Graham D, Jia H, Uchida M, Larson EL. California Hospitals Response to State and Federal Policies Related to Health Care-Associated Infections. *Policy, Politics & Nursing Practice*. 2011;12(2):73-81. doi: 10.1177/1527154411416129.
141. Meddings JA, Reichert H, Rogers MA, Saint S, Stephansky J, McMahon LF. Effect of Nonpayment for Hospital-Acquired, Catheter-Associated Urinary Tract Infection: A Statewide Analysis. *Annals of Internal Medicine*. 2012;157(5):305-312.
142. Schuller K, Probst J, Hardin J, Bennett K, Martin A. Initial Impact of Medicare's Nonpayment Policy on Catheter-Associated Urinary Tract Infections by Hospital Characteristics. *Health Policy*. 2014;115(2):165-171.
143. Gidwani R, Bhattacharya J. CMS Reimbursement Reform and the Incidence of Hospital-Acquired Pulmonary Embolism or Deep Vein Thrombosis. *Journal of General Internal Medicine*. 2015;30(5):588-596.
144. Waters TM, Daniels MJ, Bazzoli GJ, et al. Effect of Medicare's Nonpayment for Hospital-Acquired Conditions: Lessons for Future Policy. *JAMA Internal Medicine*. 2015;175(3):347-354.

145. Vaz LE, Kleinman KP, Kawai AT, et al. Impact of Medicare's Hospital-Acquired Condition Policy on Infections in Safety Net and Non-Safety Net Hospitals. *Infection Control & Hospital Epidemiology*. 2015;36(06):649-655.
146. Hoff T, Hartmann CW, Soerensen C, Wroe P, Dutta-Linn M, Lee G. Making the CMS Payment Policy for Healthcare-Associated Infections Work: Organizational Factors That Matter. *Journal of Healthcare Management*. 2011;56(5):319-335.
147. Krein SL, Kowalski CP, Hofer TP, Saint S. Preventing Hospital-Acquired Infections: A National Survey of Practices Reported by U.S. Hospitals in 2005 and 2009. *Journal of General Internal Medicine*. 2012;27(7):773-779. doi: 10.1007/s11606-011-1935-y.
148. Lee GM, Hartmann CW, Graham D, et al. Perceived Impact of the Medicare Policy to Adjust Payment for Healthcare-Associated Infections. *American Journal of Infection Control*. 2012;40(4):314-319. doi: 10.1016/j.ajic.2011.11.003.
149. McHugh M, Van Dyke K, Osei-Anto A, Haque A. Medicare's Payment Policy for Hospital-Acquired Conditions: Perspectives of Administrators From Safety Net Hospitals. *Medical Care Research and Review*. 2011;68(6):667-682. doi: 10.1177/1077558711408326.
150. Encinosa WE, Bernard DM. Hospital finances and patient safety outcomes. *Inquiry*. 2005;42(1):60-72.
151. Jha AK, Orav EJ, Li Z, Epstein AM. Concentration and quality of hospitals that care for elderly black patients. *Arch Intern Med*. 2007;167(11):1177.
152. Jha AK, Orav EJ, Epstein AM. Low-quality, high-cost hospitals, mainly in South, care for sharply higher shares of elderly black, Hispanic, and Medicaid patients. *Health Aff*. 2011;30(10):1904-1911.
153. Dimick J, Ruhter J, Sarrazin MV, Birkmeyer JD. Black Patients More Likely Than Whites To Undergo Surgery At Low-Quality Hospitals In Segregated Regions. *Health Aff*. 2013;32(6):1046-1053.
154. Institute of Medicine. *America's health care safety net: intact but endangered*. National Academies Press; 2000.
155. Goldman LE, Vittinghoff E, Dudley RA. Quality of care in hospitals with a high percent of Medicaid patients. *Med Care*. 2007;45(6):579-583.
156. Jha AK, Orav EJ, Epstein AM. The effect of financial incentives on hospitals that serve poor patients. *Ann Intern Med*. 2010;153(5):299-306.
157. Werner RM, Asch DA. The unintended consequences of publicly reporting quality information. *JAMA: the journal of the American Medical Association*. 2005;293(10):1239-1244.
158. Williams SC, Schmaltz SP, Morton DJ, Koss RG, Loeb JM. Quality of care in US hospitals as reflected by standardized measures, 2002–2004. *N Engl J Med*. 2005;353(3):255-264.
159. Conrad DA, Saver BG, Heath S. Paying physicians for quality: evidence and themes from the field. *Joint Commission Journal on Quality and Patient Safety*. 2006;32(8):443-451.

160. Dudley RA. Pay-for-performance research. *JAMA: the journal of the American Medical Association*. 2005;294(14):1821-1823.
161. Casalino LP, Elster A, Eisenberg A, Lewis E, Montgomery J, Ramos D. Will pay-for-performance and quality reporting affect health care disparities?. *Health Aff*. 2007;26(3):w405-w414.
162. Chien AT, Chin MH, Davis AM, Casalino LP. Pay for Performance, Public Reporting, and Racial Disparities in Health Care How Are Programs Being Designed?. *Medical Care Research and Review*. 2007;64(5 suppl):283S-304S.
163. Rosenthal MB, Dudley RA. Pay-for-performance: Will the latest trend improve care?. *JAMA: the journal of the American Medical Association*. 2007;297(7):740-744.
164. Epstein AM, Lee TH, Hamel MB. Paying physicians for high-quality care. *N Engl J Med*. 2004;350(4):406-410.
165. Werner RM. Does pay-for-performance steal from the poor and give to the rich?. *Ann Intern Med*. 2010;153(5):340-341.
166. Chien AT. The potential impact of performance incentive programs on racial disparities in health care. In: *Healthcare Disparities at the Crossroads with Healthcare Reform*. Springer; 2011:211-229.
167. Chin M, Chien A. Reducing racial and ethnic disparities in health care: an integral part of quality improvement scholarship. *Quality and Safety in Health Care*. 2006;15(2):79-80.
168. Ryan AM. Will Value-Based Purchasing Increase Disparities in Care?. *N Engl J Med*. 2013;369(26):2472-2474. doi: 10.1056/NEJMp1312654.
169. Burgess DJ, Fu SS, Van Ryn M. Why do providers contribute to disparities and what can be done about it?. *Journal of General Internal Medicine*. 2004;19(11):1154-1159.
170. Hasnain-Wynia R, Baker DW, Nerenz D, et al. Disparities in health care are driven by where minority patients seek care: examination of the hospital quality alliance measures. *Arch Intern Med*. 2007;167(12):1233.
171. Kressin NR. Separate but Not Equal The Consequences of Segregated Health Care. *Circulation*. 2005;112(17):2582-2584.
172. Marquis MS, Rogowski JA, Escarce JJ. Recent trends and geographic variation in the safety net. *Med Care*. 2004;42(5):408-415.
173. Werner RM, Asch DA, Polsky D. Racial profiling - The unintended consequences of coronary artery bypass graft report cards. *Circulation*. 2005;111(10):1257-1263. doi: 10.1116/01.CIR.000157729.59754.09.
174. Chaudhry B, Wang J, Wu S, et al. Systematic review: impact of health information technology on quality, efficiency, and costs of medical care. *Ann Intern Med*. 2006;144(10):742-752.
175. Chen C, Garrido T, Chock D, Okawa G, Liang L. The Kaiser Permanente Electronic Health Record: transforming and streamlining modalities of care. *Health Aff*. 2009;28(2):323-333.

176. Evans DC, Nichol WP, Perlin JB. Effect of the implementation of an enterprise-wide Electronic Health Record on productivity in the Veterans Health Administration. *Health Economics, Policy and Law*. 2006;1(02):163-169.
177. Goldzweig CL, Towfigh A, Maglione M, Shekelle PG. Costs and benefits of health information technology: new trends from the literature. *Health Aff (Millwood)*. 2009;28(2):w282-93. doi: 10.1377/hlthaff.28.2.w282; 10.1377/hlthaff.28.2.w282.
178. Hillestad R, Bigelow J, Bower A, et al. Can electronic medical record systems transform health care? Potential health benefits, savings, and costs. *Health Aff*. 2005;24(5):1103-1117.
179. Bates DW, Evans RS, Murff H, Stetson PD, Pizziferri L, Hripcsak G. Detecting adverse events using information technology. *Journal of the American Medical Informatics Association*. 2003;10(2):115-128.
180. Bates DW, Gawande AA. Improving safety with information technology. *N Engl J Med*. 2003;348(25):2526-2534.
181. Bates D, Cohen M, Leape L, Overhage JM, Shabot MM, Sheridan T. Reducing the frequency of errors in medicine using information technology. *Journal of the American Medical Informatics Association*. 2001;8(4):299-308.
182. Shapiro J. Industry preaches safety in Pittsburgh. *US News & World Report*. 2000;56.
183. Jones SS, Rudin RS, Perry T, Shekelle PG. Health Information Technology: An Updated Systematic Review With a Focus on Meaningful Use. *Annals of Internal Medicine*. 2014;160(1):48-54. doi: 10.7326/M13-1531.
184. Buntin MB, Burke MF, Hoaglin MC, Blumenthal D. The Benefits Of Health Information Technology: A Review Of The Recent Literature Shows Predominantly Positive Results. *Health Affairs*. 2011;30(3):464-471. doi: 10.1377/hlthaff.2011.0178.
185. Buising KL, Thursky KA, Robertson MB, et al. Electronic antibiotic stewardship--reduced consumption of broad-spectrum antibiotics using a computerized antimicrobial approval system in a hospital setting. *J Antimicrob Chemother*. 2008;62(3):608-616. doi: 10.1093/jac/dkn218; 10.1093/jac/dkn218.
186. Humphries TL, Carroll N, Chester EA, Magid D, Rocho B. Evaluation of an electronic critical drug interaction program coupled with active pharmacist intervention. *Ann Pharmacother*. 2007;41(12):1979-1985. doi: 10.1345/aph.1K349.
187. Rupp MT, Warholak TL. Evaluation of e-prescribing in chain community pharmacy: best-practice recommendations. *Journal of the American Pharmacists Association: JAPhA*. 2008;48(3).
188. Stone WM, Smith BE, Shaft JD, Nelson RD, Money SR. Impact of a computerized physician order-entry system. *J Am Coll Surg*. 2009;208(5):960-967.
189. DesRoches CM, Campbell EG, Rao SR, et al. Electronic health records in ambulatory care—a national survey of physicians. *N Engl J Med*. 2008;359(1):50-60.

190. Eden KB, Messina R, Li H, Osterweil P, Henderson CR, Guise J. Examining the value of electronic health records on labor and delivery. *Obstet Gynecol*. 2008;199(3):307. e1-307. e9.
191. Shachak A, Hadas-Dayagi M, Ziv A, Reis S. Primary care physicians' use of an electronic medical record system: a cognitive task analysis. *Journal of general internal medicine*. 2009;24(3):341-348.
192. Smith LB, Banner L, Lozano D, Olney CM, Friedman B. Connected care: reducing errors through automated vital signs data upload. *Comput Inform Nurs*. 2009;27(5):318-323. doi: 10.1097/NCN.0b013e3181b21d65; 10.1097/NCN.0b013e3181b21d65.
193. Evans RS, Wallace CJ, Lloyd JF, et al. Rapid identification of hospitalized patients at high risk for MRSA carriage. *Journal of the American Medical Informatics Association*. 2008;15(4):506-512.
194. Kac G, Grohs P, Durieux P, et al. Impact of electronic alerts on isolation precautions for patients with multidrug-resistant bacteria. *Arch Intern Med*. 2007;167(19):2086-2090.
195. Kho AN, Dexter PR, Warvel JS, et al. An effective computerized reminder for contact isolation of patients colonized or infected with resistant organisms. *Int J Med Inf*. 2008;77(3):194-198.
196. Parente ST, McCullough JS. Health information technology and patient safety: evidence from panel data. *Health Aff (Millwood)*. 2009;28(2):357-360. doi: 10.1377/hlthaff.28.2.357; 10.1377/hlthaff.28.2.357.
197. Gunningberg L, Fogelberg-Dahm M, Ehrenberg A. Improved quality and comprehensiveness in nursing documentation of pressure ulcers after implementing an electronic health record in hospital care. *J Clin Nurs*. 2009;18(11):1557-1564.
198. Gunningberg L, Dahm MF, Ehrenberg A. Accuracy in the recording of pressure ulcers and prevention after implementing an electronic health record in hospital care. *Qual Saf Health Care*. 2008;17(4):281-285. doi: 10.1136/qshc.2007.023341; 10.1136/qshc.2007.023341.
199. Henriksen K, Battles JB, Marks ES, et al. Clinical informatics and its usefulness for assessing risk and preventing falls and pressure ulcers in nursing home environments. 2005.
200. Willson D, Ashton C, Wingate N, et al. Computerized support of pressure ulcer prevention and treatment protocols. 1995;19:646-650.
201. McNerney JA. Reducing hospital-acquired pressure ulcer prevalence through a focused prevention program. *Adv Skin Wound Care*. 2008;21(2):75-78. doi: 10.1097/01.ASW.0000305410.58350.34; 10.1097/01.ASW.0000305410.58350.34.
202. Hunderfund ANL, Sweeney CM, Mandrekar JN, Johnson LM, Britton JW. Effect of a multidisciplinary fall risk assessment on falls among neurology inpatients. 2011;86(1):19-24.
203. Rauch K, Balascio J, Gilbert P. Excellence in action: developing and implementing a fall prevention program. *Journal for Healthcare Quality*. 2009;31(1):36-42.

204. Smith SA, Murphy ME, Huschka TR, et al. Impact of a diabetes electronic management system on the care of patients seen in a subspecialty diabetes clinic. *Diabetes Care*. 1998;21(6):972-976.
205. Balas EA, Krishna S, Kretschmer RA, Cheek TR, Lobach DF, Boren SA. Computerized knowledge management in diabetes care. *Med Care*. 2004;42(6):610-621.
206. O'Connor PJ, Crain AL, Rush WA, Sperl-Hillen JAM, Gutenkauf JJ, Duncan JE. Impact of an electronic medical record on diabetes quality of care. *The Annals of Family Medicine*. 2005;3(4):300-306.
207. Sequist TD, Gandhi TK, Karson AS, et al. A randomized trial of electronic clinical reminders to improve quality of care for diabetes and coronary artery disease. *Journal of the American Medical Informatics Association*. 2005;12(4):431-437.
208. Cebul RD, Love TE, Jain AK, Hebert CJ. Electronic health records and quality of diabetes care. *N Engl J Med*. 2011;365(9):825-833.
209. Guerra YS, Das K, Antonopoulos P, et al. Computerized physician order entry-based hyperglycemia inpatient protocol and glycemic outcomes: The CPOE-HIP study. *Endocrine Practice*. 2010;16(3):389-397.
210. Crosson JC, Ohman-Strickland PA, Hahn KA, et al. Electronic medical records and diabetes quality of care: results from a sample of family medicine practices. *The Annals of Family Medicine*. 2007;5(3):209-215.
211. Lee J, Clay B, Zelazny Z, Maynard G. Computerized Algorithms: Indication-Based Ordering: A New Paradigm for Glycemic Control in Hospitalized Inpatients. *Journal of diabetes science and technology (Online)*. 2008;2(3):349.
212. Brossette SE, Hacek DM, Gavin PJ, et al. A laboratory-based, hospital-wide, electronic marker for nosocomial infection: the future of infection control surveillance?. *Am J Clin Pathol*. 2006;125(1):34-39.
213. de Bruin JS, Seeling W, Schuh C. Data use and effectiveness in electronic surveillance of healthcare associated infections in the 21st century: a systematic review. *Journal of the American Medical Informatics Association*. 2014:amiajnl-2013-002089.
214. Tokars JI, Richards C, Andrus M, et al. The changing face of surveillance for health care-associated infections. *Clin Infect Dis*. 2004;39(9):1347-1352. doi: 10.1086/425000.
215. Shepard J, Hadhazy E, Frederick J, et al. Using electronic medical records to increase the efficiency of catheter-associated urinary tract infection surveillance for National Health and Safety Network reporting. *Am J Infect Control*. 2014;42(3):e33-e36.
216. Welden LM. Electronic Health Record: Driving Evidence-Based Catheter-Associated Urinary Tract Infections (CAUTI) Care Practices. *Online Journal of Issues in Nursing*. 2013;18(3).
217. Beeler P, Kucher N, Blaser J. Sustained impact of electronic alerts on rate of prophylaxis against venous thromboembolism. *Thromb Haemost*. 2011;106(4):734.

218. Durieux P, Nizard R, Ravaud P, Mounier N, Lepage E. A clinical decision support system for prevention of venous thromboembolism. *JAMA: the journal of the American Medical Association*. 2000;283(21):2816-2821.
219. Galanter WL, Thambi M, Rosencranz H, et al. Effects of clinical decision support on venous thromboembolism risk assessment, prophylaxis, and prevention at a university teaching hospital. *American Journal of Health-System Pharmacy*. 2010;67(15):1265-1273.
220. Haut ER, Lau BD, Kraenzlin FS, et al. Improved Prophylaxis and Decreased Rates of Preventable Harm With the Use of a Mandatory Computerized Clinical Decision Support Tool for Prophylaxis for Venous Thromboembolism in Trauma. *Archives of Surgery*. 2012;147(10):901-907.
221. Kucher N, Koo S, Quiroz R, et al. Electronic alerts to prevent venous thromboembolism among hospitalized patients. *N Engl J Med*. 2005;352(10):969-977.
222. Lecumberri R, Marqués M, Díaz-Navarraz MT, et al. Maintained effectiveness of an electronic alert system to prevent venous thromboembolism among hospitalized patients. *Thromb Haemost*. 2008;100(4):699-704.
223. Mosen D, Elliott CG, Egger MJ, et al. The effect of a computerized reminder system on the prevention of postoperative venous thromboembolism. *CHEST Journal*. 2004;125(5):1635-1641.
224. Hu HM, Tzeng HM. A hospital's adoption of information technology is associated with altered risks of hospital-acquired venous thromboembolism. *Am J Med Qual*. 2012;27(4):305-312. doi: 10.1177/1062860611426349; 10.1177/1062860611426349.
225. Jha AK, DesRoches CM, Shields AE, et al. Evidence of an emerging digital divide among hospitals that care for the poor. *Health Aff (Millwood)*. 2009;28(6):w1160-70. doi: 10.1377/hlthaff.28.6.w1160; 10.1377/hlthaff.28.6.w1160.
226. DesRoches CM, Worzala C, Joshi MS, Kralovec PD, Jha AK. Small, nonteaching, and rural hospitals continue to be slow in adopting electronic health record systems. *Health Aff (Millwood)*. 2012;31(5):1092-1099. doi: 10.1377/hlthaff.2012.0153; 10.1377/hlthaff.2012.0153.
227. Lee GM. Preventing Avoidable Infectious Complications by Adjusting Payment Project. <http://www.paicap.org/about.html>. Accessed 07/12, 2016.
228. Press Ganey. National Database for Nursing Quality Indicators. <http://www.pressganey.com/solutions/clinical-quality/nursing-quality>. Accessed 07/12, 2016.
229. Tang PC, Ralston M, Arrigotti MF, Qureshi L, Graham J. Comparison of methodologies for calculating quality measures based on administrative data versus clinical data from an electronic health record system: implications for performance measures. *J Am Med Inform Assoc*. 2007;14(1):10-15. doi: M2198 [pii].
230. Zhan C, Miller MR. Administrative data based patient safety research: a critical review. *Qual Saf Health Care*. 2003;12 Suppl 2:ii58-63.
231. Agency of Healthcare Research and Quality. State Inpatient Databases. <http://goo.gl/OFVB3Q>. Accessed 04/27, 2016.

232. Steiner C, Elixhauser A, Schnaier J. The healthcare cost and utilization project: an overview. *Effective clinical practice: ECP*. 2001;5(3):143-151.
233. Houchens RL, Elixhauser A, Romano PS. How often are potential patient safety events present on admission?. *Joint Commission Journal on Quality and Patient Safety*. 2008;34(3):154-163.
234. Coffey R, Milenkovic M, Andrews R. The case for the Present-on-Admission (POA) indicator. Rockville, MD: US Agency for Healthcare Research and Quality. 2006.
235. Centers for Medicare and Medicaid Services. Impact Files. <https://goo.gl/YQDAHV>. Accessed 04/27, 2016.
236. Centers for Medicare and Medicaid Services. Provider of Services Files. <http://goo.gl/Qg4QLW>. Accessed 04/27, 2016.
237. Dartmouth Atlas Project. The Dartmouth Atlas of Health Care. <http://goo.gl/McU7Cn>. Accessed 04/27, 2016.
238. HIMSS. HIMSS Analytics. <http://goo.gl/PbaltM>. Accessed 03/10, 2016.
239. Centers for Medicare and Medicaid Services. EHR Incentive Programs. <http://goo.gl/3ylzH1>. Accessed 04/27, 2016.
240. Centers for Medicare and Medicaid Services. Cost Reports. <http://goo.gl/qyHxms>. Accessed 04/27, 2016.
241. American Hospital Association. AHA Data Viewer. <http://goo.gl/vi7KvU>. Accessed 04/27, 2016.
242. Dharmarajan K, Hsieh AF, Lin Z, et al. Diagnoses and timing of 30-day readmissions after hospitalization for heart failure, acute myocardial infarction, or pneumonia. *JAMA*. 2013;309(4):355-363.
243. Fonarow GC, Pan W, Saver JL, et al. Comparison of 30-day mortality models for profiling hospital performance in acute ischemic stroke with vs without adjustment for stroke severity. *JAMA*. 2012;308(3):257-264.
244. Gregg EW, Li Y, Wang J, et al. Changes in diabetes-related complications in the United States, 1990–2010. *N Engl J Med*. 2014;370(16):1514-1523.
245. Centers for Medicare and Medicaid Services. Hospital-Acquired Conditions. <https://goo.gl/bVIZre>. Accessed 04/27, 2016.
246. Kassed C, Kowlessar N, Pfunter A, Parlato J, Andrews RM. The Case for the POA Indicator: Update 2011. <http://goo.gl/FdJrKK>. Accessed 02/29, 2016.
247. Centers for Medicare and Medicaid Services. Hospital-Acquired Conditions. <https://goo.gl/rOnyhX>. Accessed 02/18, 2016.
248. Rajaram R, Chung JW, Jones AT, et al. Association of the 2011 ACGME resident duty hour reform with general surgery patient outcomes and with resident examination performance. *JAMA*. 2014;312(22):2374-2384.
249. Furukawa MF, Raghu T, Shao B. Electronic medical records, nurse staffing, and nurse-sensitive patient outcomes: Evidence from California hospitals, 1998–2007. *Health Serv Res*. 2010;45(4):941-962.

250. Appari A, Eric Johnson M, Anthony DL. Meaningful Use of Electronic Health Record Systems and Process Quality of Care: Evidence from a Panel Data Analysis of US Acute-Care Hospitals. *Health Serv Res.* 2013;48(2pt1):354-375.
251. Cutler DM, Morton FS. Hospitals, market share, and consolidation. *JAMA.* 2013;310(18):1964-1970.
252. Localio AR, Berlin JA, Ten Have TR, Kimmel SE. Adjustments for center in multicenter studies: an overview. *Ann Intern Med.* 2001;135(2):112-123.
253. Angrist JD, Pischke J. Mostly harmless econometrics: An empiricist's companion. Princeton university press; 2008.
254. Scott MA, Simonoff JS, Marx BD. The SAGE handbook of multilevel modeling. Sage; 2013.
255. Clark TS, Linzer DA. Should I use fixed or random effects?. *Political Science Research and Methods.* 2015;3(02):399-408.
256. Hox JJ, Moerbeek M, van de Schoot R. Multilevel analysis: Techniques and applications. Routledge; 2010.
257. HIMSS. Electronic Medical Record Adoption Model. <https://goo.gl/A62JBS>. Accessed 04/27, 2016.
258. Jaccard J. Interaction effects in logistic regression. Vol 135. Sage; 2001.
259. Rabe-Hesketh S, Skrondal A. Multilevel and Longitudinal Modeling using Stata. STATA press; 2012.
260. McKethan A, Jha A. Designing smarter pay-for-performance programs. *JAMA.* 2014;312(24):2617-2618. doi: 10.1001/jama.2014.15398.
261. Centers for Medicare and Medicaid Services. Hospital-Acquired Condition Reduction Program. <https://goo.gl/3p5iDz>. Accessed 06/02, 2016.
262. Kahneman D, Tversky A. Prospect theory: An analysis of decision under risk. *Econometrica: Journal of the Econometric Society.* 1979:263-291.
263. Werner RM, Dudley RA. Medicare's new hospital value-based purchasing program is likely to have only a small impact on hospital payments. *Health Aff.* 2012;31(9):1932-1940.
264. Tsai TC, Jha AK, Gawande AA, Huckman RS, Bloom N, Sadun R. Hospital board and management practices are strongly related to hospital performance on clinical quality metrics. *Health Aff (Millwood).* 2015;34(8):1304-1311. doi: 10.1377/hlthaff.2014.1282 [doi].
265. Inouye SK, Brown CJ, Tinetti ME. Medicare nonpayment, hospital falls, and unintended consequences. *N Engl J Med.* 2009;360(23):2390-2393.
266. Tinetti ME. Preventing falls in elderly persons. *N Engl J Med.* 2003;348(1):42-49.
267. Oliver D, Hopper A, Seed P. Do hospital fall prevention programs work? A systematic review. *J Am Geriatr Soc.* 2000;48(12):1679-1689.

268. Gilman M, Hockenberry JM, Adams EK, Milstein AS, Wilson IB, Becker ER. The Financial Effect of Value-Based Purchasing and the Hospital Readmissions Reduction Program on Safety-Net Hospitals in 2014: A Cohort Study. *Ann Intern Med*. 2015;163(6):427-436.
269. Bilimoria KY, Chung J, Ju MH, et al. Evaluation of surveillance bias and the validity of the venous thromboembolism quality measure. *JAMA*. 2013;310(14):1482-1489.
270. Luft HS, Hunt SS, Maerki SC. The volume-outcome relationship: practice-makes-perfect or selective-referral patterns?. *Health Serv Res*. 1987;22(2):157-182.
271. Iezzoni LI. Assessing quality using administrative data. *Ann Intern Med*. 1997;127(8_Part_2):666-674.
272. Meddings J, Saint S, McMahon Jr LF. Hospital-Acquired Catheter-Associated Urinary Tract Infection: Documentation and Coding Issues May Reduce Financial Impact of Medicare's New Payment Policy. *Hospital*. 2010;31(6):627-633.
273. Silverman E, Skinner J. Medicare upcoding and hospital ownership. *J Health Econ*. 2004;23(2):369-389.
274. Donabedian A. The quality of care: How can it be assessed?. *JAMA*. 1988;260(12):1743.

APPENDIX

Table A 1: Unadjusted hospital-level Hospital-Acquired Condition rates in the medical cohort¹

Program Phase	Pre ²					Post ³				
HAC / MUR ⁴ Quartile	1	2	3	4	Total	1	2	3	4	Total
Catheter-associated bloodstream infection (BSI)										
Hospitals (inpatient stays)	35 (22,934)	39 (39,297)	41 (24,232)	38 (20,532)	153 (106,995)	34 (92,224)	38 (161,676)	39 (92,641)	37 (78,523)	148 (425,064)
Mean rate (SD) ⁵	0.05 (0.12)	0.09 (0.18)	0.06 (0.14)	0.02 (0.06)	0.05 (0.13)	0.04 (0.05)	0.06 (0.09)	0.04 (0.06)	0.03 (0.07)	0.04 (0.07)
Hospitals with at least 1 BSI (%) ⁶	28.57	43.59	17.07	10.53	24.84	55.88	65.79	46.15	32.43	50.00
Catheter-associated urinary tract infection (UTI)										
Hospitals (inpatient stays)	37 (95,097)	41 (160,420)	41 (99,581)	40 (87,422)	159 (442,520)	34 (92,224)	38 (161,676)	39 (92,641)	37 (78,523)	148 (425,064)
Mean rate (SD)	0.01 (0.02)	0.03 (0.06)	0.04 (0.11)	0.03 (0.07)	0.03 (0.07)	0.04 (0.07)	0.04 (0.05)	0.04 (0.07)	0.03 (0.05)	0.04 (0.06)
Hospitals with at least 1 UTI (%)	21.62	46.34	36.59	35.00	35.22	41.18	55.26	48.72	37.84	45.95
Falls and trauma (FAL)										
Hospitals (inpatient stays)	37 (95,097)	41 (160,420)	41 (99,581)	40 (87,422)	159 (442,520)	34 (92,224)	38 (161,676)	39 (92,641)	37 (78,523)	148 (425,064)
Mean rate (SD)	0.09 (0.14)	0.12 (0.14)	0.09 (0.09)	0.12 (0.11)	0.11 (0.12)	0.07 (0.06)	0.12 (0.19)	0.08 (0.08)	0.12 (0.11)	0.10 (0.12)
Hospitals with at least 1 FAL (%)	54.05	85.37	73.17	77.50	72.96	70.59	81.58	71.79	72.97	74.32
Glycemic control (GLU)										
Hospitals (inpatient stays)	37 (95,097)	41 (160,420)	41 (99,581)	40 (87,422)	159 (442,520)	34 (92,224)	38 (161,676)	39 (92,641)	37 (78,523)	148 (425,064)

Program Phase	Pre ²					Post ³				
HAC / MUR ⁴ Quartile	1	2	3	4	Total	1	2	3	4	Total
Mean rate (SD)	0.02 (0.05)	0.01 (0.04)	0.00 (0.02)	0.01 (0.02)	0.01 (0.03)	0.01 (0.02)	0.00 (0.01)	0.00 (0.01)	0.02 (0.07)	0.01 (0.04)
Hospitals with at least 1 GLU (%)	21.62	29.27	7.32	17.50	18.87	26.47	15.79	10.26	10.81	15.54

¹ Cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012

² Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for BSI, and from January 2005 to September 2008 for UTI, FAL, GLU.

³ Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁴ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁵ Numbers indicate mean of the HAC rate at the hospital-level with standard deviation in parentheses. The HAC rate is computed as a percentage.

⁶ Numbers indicate percentage of hospitals with at least one HAC.

Table A 2: Unadjusted hospital-level Hospital-Acquired Condition rates in the surgical cohort¹

Program Phase	Pre ²					Post ³				
HAC / MUR ⁴ Quartile	1	2	3	4	Total	1	2	3	4	Total
n	32	40	41	40	153	31	37	39	34	141
Catheter-associated bloodstream infection (BSI)										
Hospitals (inpatient stays)	30 (4,858)	39 (11,379)	41 (9,691)	36 (6,371)	146 (32,299)	31 (19,154)	37 (51,812)	39 (44,840)	34 (25,521)	141 (141,327)
Mean rate (SD) ⁵	0.11 (0.36)	0.11 (0.36)	0.01 (0.05)	0.01 (0.07)	0.06 (0.25)	0.01 (0.04)	0.04 (0.07)	0.04 (0.13)	0.01 (0.04)	0.03 (0.08)
Hospitals with at least 1 BSI (%) ⁶	16.67	17.95	4.88	5.56	10.96	12.90	37.84	17.95	8.82	19.86
Catheter-associated urinary tract infection (UTI)										
Hospitals (inpatient stays)	32 (19,996)	40 (44,058)	41 (34,381)	40 (24,951)	153 (123,386)	31 (19,154)	37 (51,812)	39 (44,840)	34 (25,521)	141 (141,327)
Mean rate (SD)	0.01 (0.04)	0.02 (0.08)	0.06 (0.28)	0.03 (0.12)	0.03 (0.16)	0.03 (0.07)	0.01 (0.04)	0.07 (0.17)	0.02 (0.06)	0.03 (0.10)
Hospitals with at least 1 UTI (%)	9.38	15.00	12.20	12.50	12.42	16.13	18.92	25.64	14.71	19.15
Falls and trauma (FAL)										
Hospitals (inpatient stays)	32 (19,996)	40 (44,058)	41 (34,381)	40 (24,951)	153 (123,386)	31 (19,154)	37 (51,812)	39 (44,840)	34 (25,521)	141 (141,327)
Mean rate (SD)	0.47 (0.74)	0.50 (0.82)	0.30 (0.45)	0.59 (0.72)	0.46 (0.69)	0.23 (0.41)	0.31 (0.41)	0.36 (0.61)	0.26 (0.58)	0.30 (0.51)
Hospitals with at least 1 FAL (%)	56.25	72.50	56.10	72.50	64.71	45.16	81.08	74.36	52.94	64.54
Glycemic control (GLU)										
Hospitals (inpatient stays)	32 (19,996)	40 (44,058)	41 (34,381)	40 (24,951)	153 (123,386)	31 (19,154)	37 (51,812)	39 (44,840)	34 (25,521)	141 (141,327)

Program Phase	Pre ²					Post ³				
HAC / MUR ⁴ Quartile	1	2	3	4	Total	1	2	3	4	Total
Mean rate (SD)	0.07 (0.37)	0.01 (0.06)	0.00 (0.01)	0.00 (0.00)	0.02 (0.17)	0.00 (0.00)	0.01 (0.03)	0.00 (0.00)	0.00 (0.00)	0.00 (0.01)
Hospitals with at least 1 GLU (%)	9.38	10.00	2.44	2.50	5.88	0.00	13.51	0.00	0.00	3.55

¹ Cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

² Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for BSI, and from January 2005 to September 2008 for UTI, FAL, GLU, CabgSSI, SpineSSI, HipDVT, KneeDVT.

³ Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁴ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁵ Numbers indicate mean of the HAC rate at the hospital-level with standard deviation in parentheses. The HAC rate is computed as a percentage.

⁶ Range of the hospital-level HAC rate.

⁶ Numbers indicate percentage of hospitals with at least one HAC.

Table A 3: Multivariate odds ratios for the Program effects on the secondary outcomes¹ of the medical and surgical cohorts

Outcome	BSI	UTI	FRA	GLU
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Medical Cohort²				
Program⁴ x MUR⁵: Interaction effects				
Post- x MUR Quartile 2	1.18 [0.65,2.14]	0.52 [0.23,1.22]	0.86 [0.42,1.77]	0.39 [0.14,1.09]
Post- x MUR Quartile 3	1.45 [0.67,3.15]	0.30* [0.12,0.75]	1.30 [0.67,2.53]	1.74 [0.43,7.04]
Post- x MUR Quartile 4	1.84 [0.63,5.37]	0.46 [0.13,1.66]	1.33 [0.68,2.59]	0.84 [0.25,2.88]
p-value for joint test of interaction	0.61	0.09	0.43	0.19
Model parameters				
Inpatient stays	524,219	859,757	867,078	806,726
Hospitals	152	158	159	158
Random effects	0.61	0.81	0.43	0.86
Intraclass correlation coefficient	0.10	0.17	0.05	0.18
Chi-square for likelihood ratio test	33.13	99.20	71.51	14.98
Surgical Cohort³				
Program x MUR: Interaction effects				
Post- x MUR Quartile 2	3.97 [0.96,16.40]	0.60 [0.13,2.73]	1.04 [0.63,1.69]	1.00 [1.00,1.00]
Post- x MUR Quartile 3	7.40* [1.23,44.55]	0.98 [0.23,4.22]	1.25 [0.72,2.20]	1.00 [1.00,1.00]
Post- x MUR Quartile 4	8.75** [1.74,44.03]	1.08 [0.18,6.46]	0.91 [0.52,1.58]	1.00 [1.00,1.00]
p-value for joint test of interaction	0.04	0.84	0.74	-- ⁶
Model parameters				
Inpatient stays	156,230	254,808	264,155	77,210
Hospitals	120	150	153	70
Random effects	0.00	0.92	0.43	0.00
Intraclass correlation coefficient	0.00	0.20	0.05	0.00
Chi-square for likelihood ratio test	0.00	16.95	30.30	0.00

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: BSI: Catheter-associated bloodstream infection; UTI: Catheter-associated urinary tract infection; FRA: Falls and trauma; GLU: Poor glucose control; OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group; MUR: Medicare Utilization Ratio

¹ Secondary outcomes: Catheter-associated bloodstream infections (BSI), Catheter-associated urinary tract infections (UTI), falls and trauma (FAL), and poor glucose control (GLU)

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012.

³ Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

⁴ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for BSI, and January 2005 to September 2008 for UTI, FAL and GLU. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁵ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁶ Interaction term not testable.

Table A 4: Multivariate odds ratios for secondary outcomes¹ among hospitals with differential operating profit margins

Outcome	BSI	UTI	FRA	GLU
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Medical Cohort²				
Program⁴ x MUR⁵ quartile x Operating margin: 3-way interaction				
Post- # MUR Q2 # Positive margin	0.91 [0.16,4.99]	6.38 [0.90,45.43]	2.41* [1.05,5.50]	1.25 [0.30,5.29]
Post- # MUR Q3 # Positive margin	1.65 [0.20,13.69]	1.95 [0.30,12.82]	1.61 [0.66,3.92]	14.42** [1.90,109.41]
Post- # MUR Q4 # Positive margin	3.67 [0.60,22.59]	4.91 [0.46,52.14]	1.01 [0.41,2.52]	0.30 [0.02,3.96]
p-value for joint test of interaction	0.50	0.18	0.09	0.03
Model parameters				
Inpatient stays	522,039	850,409	864,735	806,726
Hospitals	151	157	159	158
Random effects	0.59	0.82	0.43	0.79
Intraclass correlation coefficient	0.09	0.17	0.05	0.16
Chi-square for likelihood ratio test	30.05	95.23	74.5	11.83
Surgical Cohort³				
Program x MUR quartile x Operating margin: 3-way interaction				
Post- # MUR Q2 # Positive margin	11.47* [1.31,100.61]	1.00 [1.00,1.00]	0.50 [0.13,1.85]	1.00 [1.00,1.00]
Post- # MUR Q3 # Positive margin	1.00 [1.00,1.00]	9.37* [1.07,82.02]	0.47 [0.12,1.91]	1.00 [1.00,1.00]
Post- # MUR Q4 # Positive margin	1.00 [1.00,1.00]	23.59 [0.49,1132.81]	0.70 [0.14,3.45]	1.00 [1.00,1.00]
p-value for joint test of interaction	0.03	0.07	0.70	-- ⁶
Model parameters				
Inpatient stays	152,975	245,650	261,760	68,536
Hospitals	118	150	153	63
Random effects	0.00	0.86	0.41	0.00
Intraclass correlation coefficient	0.00	0.18	0.05	0.00
Chi-square for likelihood ratio test	0.00	13.44	24.40	0.00

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: BSI: Catheter-associated bloodstream infection; UTI: Catheter-associated urinary tract infection; FRA: Falls and trauma; GLU: Poor glucose control; OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group; MUR: Medicare Utilization Ratio

¹ Secondary outcomes: Catheter-associated bloodstream infections (BSI), Catheter-associated urinary tract infections (UTI), falls and trauma (FAL), and poor glucose control (GLU)

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012.

³ Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

⁴ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for BSI, and January 2005 to September 2008 for UTI, FAL and GLU. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁵ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁶ Interaction term not testable.

Table A 5: Multivariate odds ratios for secondary outcomes¹ among hospitals with differential disproportionate patient percent

Outcome	BSI	UTI	FRA	GLU
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Medical Cohort²				
Program⁴ x MUR quartile⁵ x Disproportionate Patient Percent (DPP): 3-way interaction				
Post- # MUR Q2 # Above median	2.78 [0.40,19.31]	0.24 [0.05,1.28]	1.91 [0.43,8.54]	0.68 [0.07,6.90]
Post- # MUR Q3 # Above median	1.00 [1.00,1.00]	0.47 [0.08,2.68]	3.16 [0.67,14.98]	1.00 [1.00,1.00]
Post- # MUR Q4 # Above median	1.00 [1.00,1.00]	1.00 [1.00,1.00]	5.73* [1.18,27.74]	1.00 [1.00,1.00]
p-value for joint test of interaction	0.30	0.25	0.10	0.75
Model parameters				
Inpatient stays	523,298	857,476	867,078	794,498
Hospitals	152	158	159	157
Random effects	0.58	0.81	0.43	0.86
Intraclass correlation coefficient	0.09	0.17	0.05	0.18
Chi-square for likelihood ratio test	24.32	88.46	74.56	16.00
Surgical Cohort³				
Program x MUR quartile x DPP: 3-way interaction				
Post- # MUR Q2 # Above median	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.03 [0.11,9.62]	1.00 [1.00,1.00]
Post- # MUR Q3 # Above median	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.03 [0.11,9.78]	1.00 [1.00,1.00]
Post- # MUR Q4 # Above median	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
p-value for joint test of interaction	-- ⁶	--	1.00	--
Model parameters				
Inpatient stays	155,315	228,956	263,717	64,344
Hospitals	119	142	153	68
Random effects	0.00	1.00	0.42	0.00
Intraclass correlation coefficient	0.00	0.23	0.05	0.00
Chi-square for likelihood ratio test	0.00	13.88	27.69	0.00

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: BSI: Catheter-associated bloodstream infection; UTI: Catheter-associated urinary tract infection; FRA: Falls and trauma; GLU: Poor glucose control; OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group; MUR: Medicare Utilization Ratio

¹ Secondary outcomes: Catheter-associated bloodstream infections (BSI), Catheter-associated urinary tract infections (UTI), falls and trauma (FAL), and poor glucose control (GLU)

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012.

³ Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

⁴ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for BSI, and January 2005 to September 2008 for UTI, FAL and GLU. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁵ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁶ Interaction term not testable.

Table A 6: Multivariate odds ratios for secondary outcomes¹ among hospitals with differential Electronic Health Record (EHR) capabilities

Outcome	BSI	UTI	FRA	GLU
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
Medical Cohort²				
Program⁴ x MUR⁵ quartile x Electronic Medical Record Adoption Model (EMRAM) Stage: 3-way interaction				
Post- # MUR Q2 # Stage 2	1.00 [1.00,1.00]	1.00 [1.00,1.00]	0.13* [0.02,0.70]	1.00 [1.00,1.00]
Post- # MUR Q2 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q3 # Stage 2	0.41 [0.02,10.05]	0.14** [0.04,0.58]	0.30 [0.05,1.64]	1.00 [1.00,1.00]
Post- # MUR Q3 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q4 # Stage 2	1.00 [1.00,1.00]	1.00 [1.00,1.00]	0.63 [0.09,4.15]	1.00 [1.00,1.00]
Post- # MUR Q4 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
p-value for joint test of interaction	0.58	0.01	0.03	-- ⁶
Model parameters				
Inpatient stays	515,106	844,692	858,434	746,446
Hospitals	150	156	158	156
Random effects	0.59	0.80	0.44	0.85
Intraclass correlation coefficient	0.10	0.16	0.06	0.18
Chi-square for likelihood ratio test	30.57	85.39	69.91	14.81
Surgical Cohort³				
Program x MUR quartile x Electronic Medical Record Adoption Model (EMRAM) Stage: 3-way interaction				
Post- # MUR Q2 # Stage 2	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.70 [0.13,22.34]	1.00 [1.00,1.00]
Post- # MUR Q2 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q3 # Stage 2	1.00 [1.00,1.00]	1.00 [1.00,1.00]	0.78 [0.15,4.09]	1.00 [1.00,1.00]
Post- # MUR Q3 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q4 # Stage 2	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]
Post- # MUR Q4 # Stage 3+	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]	1.00 [1.00,1.00]

Outcome	BSI	UTI	FRA	GLU
	OR [95% CI]	OR [95% CI]	OR [95% CI]	OR [95% CI]
p-value for joint test of interaction	--	--	0.78	--
Model parameters				
Inpatient stays	148,755	245,993	261,245	72,647
Hospitals	117	148	151	69
Random effects	0.00	0.80	0.43	0.00
Intraclass correlation coefficient	0.00	0.16	0.05	0.00
Chi-square for likelihood ratio test	0.00	10.03	28.88	0.00

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Abbreviations: BSI: Catheter-associated bloodstream infection; UTI: Catheter-associated urinary tract infection; FRA: Falls and trauma; GLU: Poor glucose control; OR: Odds Ratio; CI: Confidence Interval; Ref: Reference group; MUR: Medicare Utilization Ratio

¹ Secondary outcomes: Catheter-associated bloodstream infections (BSI), Catheter-associated urinary tract infections (UTI), falls and trauma (FAL), and poor glucose control (GLU)

² Medical cohort comprises of inpatient stays for elderly Medicare beneficiaries with primary diagnosis of acute myocardial infarction, congestive heart failure, pneumonia or stroke discharged from New York hospitals from 2005-2012.

³ Surgical cohort comprises of inpatient stays for elderly Medicare beneficiaries undergoing Coronary Artery Bypass Graft, Spine surgeries, Hip or Knee replacements in New York hospitals from 2005-2012.

⁴ Pre-phase includes inpatient stays with discharges from October 2007 to September 2008 for BSI, and January 2005 to September 2008 for UTI, FAL and GLU. Post-phase includes inpatient stays with discharges from October 2008 to December 2012.

⁵ Medicare Utilization Ratio is the percent of inpatient days in a hospital that are attributed to Medicare.

⁶ 3-way interaction term not testable.